


# High Energy Jets

Andreas Maier



# The High Energy Limit

## Multi-Regge Kinematics




$$y_1 < y_2 < \dots < y_{n-1} < y_n$$

High energy limit:

- All rapidity gaps large, all transverse momenta similar
- All outgoing invariant masses  $\gg$  all  $t$ -channel momenta

# Example: Higgs Boson Production With Jets

Weak Boson Fusion



Gluon Fusion



- Known at NNLO,  $\sim 5\%$  correction


[Cacciari, Dreyer, Karlberg, Salam, Zanderighi 2015; Cruz-Martinez, Gehrmann, Glover, Huss 2018]

- Probes couplings to  $W, Z$  bosons
- Widely separated jets, WBF cuts:


$$y_{j_1 j_2} > 2.8, \quad m_{j_1 j_2} > 400 \text{ GeV}$$

# Example: Higgs Boson Production With Jets

Weak Boson Fusion



Gluon Fusion




- Known at LO for  $H + 2, 3$  jets,  $m_t \rightarrow \infty$  is off by  $\sim 2\%$

[Greiner, Höche, Luisoni, Schönherr, Winter 2016]

- $H + 2, 3$  jets known at NLO for  $m_t \rightarrow \infty$ .


[Cullen et al. 2013]



plot from [Greiner, Höche, Luisoni, Schönherr, Winter, Yundin 2015]


# The High Energy Limit

FKL configurations [Fadin, Kuraev, Lipatov 1975–1977]:



- Maximum number of possible  $t$ -channel gluon exchanges
- Dominant contribution in high-energy limit


Non-FKL configurations:



- Less possible  $t$ -channel gluon exchanges
- Exponentially suppressed at large rapidities

# The High Energy Limit


## Scaling Behaviour



[Andersen, Hapola, Maier, Smillie 2017]

# High Energy Jets (HEJ) resummation

**FKL Matrix element** [Andersen, Del Duca, Smillie, White 2008–2010]



Increasing rapidity

$$\begin{aligned} \left| \mathcal{M}_{f_1 f_2 \rightarrow f_1 \cdot g \cdot H \cdot g \cdot f_2}^{\text{HEJ}} \right|^2 = \\ \frac{1}{4(N_C^2 - 1)} \| j_\mu V_H^{\mu\nu} j_\nu \|^2 \cdot \left( \frac{1}{t_j t_{j+1}} \right) \\ \cdot \left( g_s^2 K_{f_1}(p_1^-, p_a^-) \frac{1}{t_1} \right) \\ \cdot \left( g_s^2 K_{f_2}(p_n^+, p_b^+) \frac{1}{t_n} \right) \\ \cdot \prod_{\substack{1 \leq k < j \\ j+1 \leq k < n}} \frac{-g_s^2 C_A}{t_k t_{k+1}} V_L(q_k, q_{k+1})^2 \\ \cdot \prod_{i=1}^{n-1} \exp \left[ \omega^0(q_{i\perp})(y_{i+1} - y_i) \right]. \end{aligned}$$


$$K_q = C_F, \quad K_g(p_1^-, p_a^-) \xrightarrow{p_1^- \rightarrow p_a^-} C_A,$$

$$\omega^0(q_{i\perp}) = -C_A \frac{\alpha_s}{\pi} \log \frac{q_{i\perp}^2}{\lambda^2}.$$

# High Energy Jets (HEJ) resummation


## Unordered Emissions

Forward or backward gluon emission:





[Andersen, Hapola, Maier, Smillie 2017]

Forward or backward Higgs boson emission:




# High Energy Jets (HEJ) approximation

## Comparison to fixed order




# High Energy Jets (HEJ) resummation

Matching to leading order



Fixed-order FKL event  
MadGraph, Sherpa, ...  
 $\sim |\mathcal{M}_{\text{LO}}|^2$



Resummation events  
Keep Higgs + jet rapidities, shift jet  $p_\perp$   
 $\sim |\mathcal{M}_{\text{HEJ}}|^2$


$$\text{Final resummation event weight} \sim \frac{|\mathcal{M}_{\text{LO}}|^2 |\mathcal{M}_{\text{HEJ}}|^2}{|\mathcal{M}_{\text{HEJ, LO}}|^2}$$

[Andersen, Hapola, Heil, Maier, Smillie 2018]

# High Energy Jets (HEJ) resummation



Matching to parton shower [Andersen, Brooks, Lönnblad 2018]

Add extra soft and collinear radiation



Subtract double counting with *modified splitting function*



$$P^{\text{subtracted}} = P^{\text{PYTHIA}} - \frac{1}{2} \frac{1}{16\pi^2} \frac{|\mathcal{M}_{n+1}^{\text{HEJ}}|^2}{|\mathcal{M}_n^{\text{HEJ}}|^2}$$



# Results for Higgs+jets

[Andersen, Hapola, Heil, Maier, Smillie 2017–2018]

# Invariant mass distribution





$$\mu = H_T/2$$


$$\mu = \max(m_{j_1,j_2}, m_h)$$

$\sigma_{\text{HEJ}}$  rescaled to  $\sigma_{\text{NLO}}$


# Jet multiplicity



# Higgs transverse momentum distribution




inclusive



WBF cuts




# Higgs transverse momentum distribution

## Quark mass effects




# Jet veto efficiency


[Rainwater, Szalapski, Zeppenfeld 1996]



# Angular separation



inclusive



WBF cuts

Angular correlation  $\Rightarrow$  CP properties of  $ggH$  coupling

# Conclusion

- High Energy Jets (HEJ) provides all-order resummation for large rapidity spans
- Relevant for Higgs + jets in gluon fusion:  
Predictions differ significantly from NLO
- Other processes:  $W, Z/\gamma + \text{jets}$

[Andersen, Hapola, Smillie 2012; Andersen, Medley, Smillie 2016]

Work in progress:

- Quark mass effects in Higgs + jets
- Fixed-order matching at and beyond LO
- Subleading corrections
- Combination with parton showers
- Further processes

# Backup

# Comparison for $W + \text{jets}$

[arXiv:1703.04362]

