

Precision in EFT studies for top and Higgs physics

Eleni Vryonidou
CERN TH




University of Vienna
22/10/19

Outline


- Introduction to the EFT
- EFT in top quark physics
 - Precision calculations in the EFT
 - Towards global fits in the top sector
- EFT in the top-Higgs sector
 - Top loops in the EFT

LHC: the story so far

Higgs discovery



Rediscovering the SM




Good agreement with the SM predictions

How to look for new physics?

Model-dependent

SUSY, 2HDM...

New particles




Resonance peaks

Model-Independent

simplified models, EFT


New Interactions
of SM particles

anomalous couplings, EFT




Deviations in tails


SMEFT: What is it all about?




SMEFT: What is it all about?




SMEFT: What is it all about?




SMEFT: What is it all about?




SMEFT: What is it all about?




SMEFT: What is it all about?




SMEFT: What is it all about?



SMEFT: What is it all about?



SMEFT basics



New Interactions of SM particles

$$\mathcal{L}_{\text{Eff}} = \mathcal{L}_{\text{SM}} + \sum_i \frac{C_i^{(6)} O_i^{(6)}}{\Lambda^2} + \mathcal{O}(\Lambda^{-4})$$

Buchmuller, Wyler Nucl.Phys. B268 (1986) 621-653

Grzadkowski et al arXiv:1008.4884

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_φ	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{WB}}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_p \gamma^\mu d_r)$

$(\bar{L}L)(\bar{L}L)$		$(\bar{R}R)(\bar{R}R)$		$(\bar{L}L)(\bar{R}R)$	
Q_{ll}	$(\bar{l}_p \gamma_\mu l_r)(\bar{l}_s \gamma^\mu l_t)$	Q_{ee}	$(\bar{e}_p \gamma_\mu e_r)(\bar{e}_s \gamma^\mu e_t)$	Q_{le}	$(\bar{l}_p \gamma_\mu l_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{qq}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{uu}	$(\bar{u}_p \gamma_\mu u_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{lu}	$(\bar{l}_p \gamma_\mu l_r)(\bar{u}_s \gamma^\mu u_t)$
$Q_{qq}^{(3)}$	$(\bar{q}_p \gamma_\mu \tau^I q_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{dd}	$(\bar{d}_p \gamma_\mu d_r)(\bar{d}_s \gamma^\mu d_t)$	Q_{ld}	$(\bar{l}_p \gamma_\mu l_r)(\bar{d}_s \gamma^\mu d_t)$
$Q_{lq}^{(1)}$	$(\bar{l}_p \gamma_\mu l_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{eu}	$(\bar{e}_p \gamma_\mu e_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{qe}	$(\bar{q}_p \gamma_\mu q_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{lq}^{(3)}$	$(\bar{l}_p \gamma_\mu \tau^I l_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{ed}	$(\bar{e}_p \gamma_\mu e_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{u}_s \gamma^\mu u_t)$
		$Q_{ud}^{(1)}$	$(\bar{u}_p \gamma_\mu u_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(8)}$	$(\bar{q}_p \gamma_\mu T^A q_r)(\bar{u}_s \gamma^\mu T^A u_t)$
		$Q_{ud}^{(8)}$	$(\bar{u}_p \gamma_\mu T^A u_r)(\bar{d}_s \gamma^\mu T^A d_t)$	$Q_{qd}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{d}_s \gamma^\mu d_t)$
				$Q_{qd}^{(8)}$	$(\bar{q}_p \gamma_\mu T^A q_r)(\bar{d}_s \gamma^\mu T^A d_t)$
$(\bar{L}R)(\bar{R}L)$ and $(\bar{L}R)(\bar{L}R)$		B -violating			
Q_{ledq}	$(\bar{l}_p e_r)(\bar{d}_s q_t^i)$	Q_{duq}	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{ijk} [(d_p^\alpha)^T C u_r^\beta] [(q_s^\gamma)^T C l_t^k]$		
$Q_{quqd}^{(1)}$	$(\bar{q}_p^j u_r) \varepsilon_{jk} (\bar{q}_s^k d_t)$	Q_{qqq}	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{ijk} [(q_p^\alpha)^T C q_r^\beta] [(u_s^\gamma)^T C e_t]$		
$Q_{quqd}^{(8)}$	$(\bar{q}_p^j T^A u_r) \varepsilon_{jk} (\bar{q}_s^k T^A d_t)$	$Q_{qqq}^{(1)}$	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{ijk} \varepsilon_{mnn} [(q_p^\alpha)^T C q_r^\beta] [(q_s^\gamma)^T C l_t^n]$		
$Q_{lequ}^{(1)}$	$(\bar{l}_p e_r) \varepsilon_{jk} (\bar{q}_s^k u_t)$	$Q_{qqq}^{(3)}$	$\varepsilon^{\alpha\beta\gamma} (\tau^I \varepsilon)_{jk} (\tau^I \varepsilon)_{mn} [(q_p^\alpha)^T C q_r^\beta] [(q_s^\gamma)^T C l_t^n]$		
$Q_{lequ}^{(3)}$	$(\bar{l}_p \sigma_{\mu\nu} e_r) \varepsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t)$	Q_{duu}	$\varepsilon^{\alpha\beta\gamma} [(d_p^\alpha)^T C u_r^\beta] [(u_s^\gamma)^T C e_t]$		

Outline

- Introduction to the EFT
- EFT in top quark physics
 - Precision calculations in the EFT
 - Towards global fits in the top sector
- EFT in the top-Higgs sector
 - Top loops in the EFT

EFT for top quark interactions

SMEFT

vs

Anomalous couplings

$$O_{\varphi Q}^{(3)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu^I \varphi \right) (\bar{Q}\gamma^\mu \tau^I Q)$$

$$O_{\varphi Q}^{(1)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{Q}\gamma^\mu Q)$$

$$O_{\varphi t} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{t}\gamma^\mu t)$$

$$O_{tW} = y_t g_w (\bar{Q}\sigma^{\mu\nu} \tau^I t) \tilde{\varphi} W_{\mu\nu}^I$$

$$O_{tB} = y_t g_Y (\bar{Q}\sigma^{\mu\nu} t) \tilde{\varphi} B_{\mu\nu}$$

$$\mathcal{L}_{ttZ} = e\bar{u}(p_t) \left[\gamma^\mu (C_{1,V}^Z + \gamma_5 C_{1,A}^Z) + \frac{i\sigma^{\mu\nu} q_\nu}{m_Z} (C_{2,V}^Z + i\gamma_5 C_{2,A}^Z) \right] v(p_{\bar{t}}) Z_\mu$$

- SMEFT:
 - Gauge invariant ✓
 - Higher-order corrections: renormalisable order by order in $1/\Lambda$ ✓


$$\mathcal{O}(\alpha_s) + \mathcal{O}\left(\frac{1}{\Lambda^2}\right) + \mathcal{O}\left(\frac{\alpha_s}{\Lambda^2}\right) + \dots$$

- Complete description-respecting SM symmetries ✓
- Model Independent ✓


SMEFT in processes with tops

Rich phenomenology:



pair production








single



associated production



top loops



connection to Higgs physics

Top-quark operators & how to look for them

$$O_{\varphi Q}^{(3)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu^I \varphi \right) (\bar{Q}\gamma^\mu \tau^I Q)$$

$$O_{\varphi Q}^{(1)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{Q}\gamma^\mu Q)$$

$$O_{\varphi t} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{t}\gamma^\mu t)$$

$$O_{tW} = y_t g_w (\bar{Q}\sigma^{\mu\nu} \tau^I t) \tilde{\varphi} W_{\mu\nu}^I$$

$$O_{tB} = y_t g_Y (\bar{Q}\sigma^{\mu\nu} t) \tilde{\varphi} B_{\mu\nu}$$

$$O_{tG} = y_t g_s (\bar{Q}\sigma^{\mu\nu} T^A t) \tilde{\varphi} G_{\mu\nu}^A,$$


$$O_{t\phi} = y_t^3 \left(\phi^\dagger \phi \right) (\bar{Q}t) \tilde{\phi}$$

see for example: Aguilar-Saavedra (arXiv:0811.3842)

Zhang and Willenbrock (arXiv:1008.3869)

+four-fermion operators

+non-top operators (mixing)



Top-quark operators & how to look for them

$$O_{\varphi Q}^{(3)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu^I \varphi \right) (\bar{Q}\gamma^\mu \tau^I Q)$$

$$O_{\varphi Q}^{(1)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{Q}\gamma^\mu Q)$$

$$O_{\varphi t} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{t}\gamma^\mu t)$$

$$O_{tW} = y_t g_w (\bar{Q}\sigma^{\mu\nu} \tau^I t) \tilde{\varphi} W_{\mu\nu}^I$$

$$O_{tB} = y_t g_Y (\bar{Q}\sigma^{\mu\nu} t) \tilde{\varphi} B_{\mu\nu}$$

$$O_{tG} = y_t g_s (\bar{Q}\sigma^{\mu\nu} T^A t) \tilde{\varphi} G_{\mu\nu}^A,$$


$$O_{t\phi} = y_t^3 \left(\phi^\dagger \phi \right) (\bar{Q}t) \tilde{\phi}$$

see for example: Aguilar-Saavedra (arXiv:0811.3842)

Zhang and Willenbrock (arXiv:1008.3869)

+four-fermion operators

+non-top operators (mixing)



Top-quark operators & how to look for them

$$O_{\varphi Q}^{(3)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu^I \varphi \right) (\bar{Q}\gamma^\mu \tau^I Q)$$

$$O_{\varphi Q}^{(1)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{Q}\gamma^\mu Q)$$

$$O_{\varphi t} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{t}\gamma^\mu t)$$

$$O_{tW} = y_t g_w (\bar{Q}\sigma^{\mu\nu} \tau^I t) \tilde{\varphi} W_{\mu\nu}^I$$

$$O_{tB} = y_t g_Y (\bar{Q}\sigma^{\mu\nu} t) \tilde{\varphi} B_{\mu\nu}$$

$$O_{tG} = y_t g_s (\bar{Q}\sigma^{\mu\nu} T^A t) \tilde{\varphi} G_{\mu\nu}^A,$$


$$O_{t\phi} = y_t^3 \left(\phi^\dagger \phi \right) (\bar{Q}t) \tilde{\phi}$$

see for example: Aguilar-Saavedra (arXiv:0811.3842)

Zhang and Willenbrock (arXiv:1008.3869)

+four-fermion operators

+non-top operators (mixing)



Top-quark operators & how to look for them

$$O_{\varphi Q}^{(3)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu^I \varphi \right) (\bar{Q}\gamma^\mu \tau^I Q)$$

$$O_{\varphi Q}^{(1)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{Q}\gamma^\mu Q)$$

$$O_{\varphi t} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{t}\gamma^\mu t)$$

$$O_{tW} = y_t g_w (\bar{Q}\sigma^{\mu\nu} \tau^I t) \tilde{\varphi} W_{\mu\nu}^I$$

$$O_{tB} = y_t g_Y (\bar{Q}\sigma^{\mu\nu} t) \tilde{\varphi} B_{\mu\nu}$$

$$O_{tG} = y_t g_s (\bar{Q}\sigma^{\mu\nu} T^A t) \tilde{\varphi} G_{\mu\nu}^A,$$


$$O_{t\phi} = y_t^3 \left(\phi^\dagger \phi \right) (\bar{Q}t) \tilde{\phi}$$

see for example: Aguilar-Saavedra (arXiv:0811.3842)

Zhang and Willenbrock (arXiv:1008.3869)

+four-fermion operators

+non-top operators (mixing)



Top-quark operators & how to look for them

$$O_{\varphi Q}^{(3)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu^I \varphi \right) (\bar{Q}\gamma^\mu \tau^I Q)$$

$$O_{\varphi Q}^{(1)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{Q}\gamma^\mu Q)$$

$$O_{\varphi t} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{t}\gamma^\mu t)$$

$$O_{tW} = y_t g_w (\bar{Q}\sigma^{\mu\nu} \tau^I t) \tilde{\varphi} W_{\mu\nu}^I$$

$$O_{tB} = y_t g_Y (\bar{Q}\sigma^{\mu\nu} t) \tilde{\varphi} B_{\mu\nu}$$

$$O_{tG} = y_t g_s (\bar{Q}\sigma^{\mu\nu} T^A t) \tilde{\varphi} G_{\mu\nu}^A,$$


$$O_{t\phi} = y_t^3 \left(\phi^\dagger \phi \right) (\bar{Q}t) \tilde{\phi}$$

see for example: Aguilar-Saavedra (arXiv:0811.3842)

Zhang and Willenbrock (arXiv:1008.3869)

+four-fermion operators

+non-top operators (mixing)



Top-quark operators & how to look for them

$$O_{\varphi Q}^{(3)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu^I \varphi \right) (\bar{Q}\gamma^\mu \tau^I Q)$$

$$O_{\varphi Q}^{(1)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{Q}\gamma^\mu Q)$$

$$O_{\varphi t} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{t}\gamma^\mu t)$$

$$O_{tW} = y_t g_w (\bar{Q}\sigma^{\mu\nu} \tau^I t) \tilde{\varphi} W_{\mu\nu}^I$$

$$O_{tB} = y_t g_Y (\bar{Q}\sigma^{\mu\nu} t) \tilde{\varphi} B_{\mu\nu}$$

$$O_{tG} = y_t g_s (\bar{Q}\sigma^{\mu\nu} T^A t) \tilde{\varphi} G_{\mu\nu}^A$$


$$O_{t\phi} = y_t^3 \left(\phi^\dagger \phi \right) (\bar{Q}t) \tilde{\phi}$$

see for example: Aguilar-Saavedra (arXiv:0811.3842)

Zhang and Willenbrock (arXiv:1008.3869)

+four-fermion operators

+non-top operators (mixing)



Top-quark operators & how to look for them

$$O_{\varphi Q}^{(3)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu^I \varphi \right) (\bar{Q}\gamma^\mu \tau^I Q)$$

$$O_{\varphi Q}^{(1)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{Q}\gamma^\mu Q)$$

$$O_{\varphi t} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{t}\gamma^\mu t)$$

$$O_{tW} = y_t g_w (\bar{Q}\sigma^{\mu\nu} \tau^I t) \tilde{\varphi} W_{\mu\nu}^I$$

$$O_{tB} = y_t g_Y (\bar{Q}\sigma^{\mu\nu} t) \tilde{\varphi} B_{\mu\nu}$$

$$O_{tG} = y_t g_s (\bar{Q}\sigma^{\mu\nu} T^A t) \tilde{\varphi} G_{\mu\nu}^A,$$


$$O_{t\phi} = y_t^3 \left(\phi^\dagger \phi \right) (\bar{Q}t) \tilde{\phi}$$

see for example: Aguilar-Saavedra (arXiv:0811.3842)

Zhang and Willenbrock (arXiv:1008.3869)

+four-fermion operators

+non-top operators (mixing)



Top-quark operators & how to look for them

$$O_{\varphi Q}^{(3)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu^I \varphi \right) (\bar{Q}\gamma^\mu \tau^I Q)$$

$$O_{\varphi Q}^{(1)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{Q}\gamma^\mu Q)$$

$$O_{\varphi t} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{t}\gamma^\mu t)$$

$$O_{tW} = y_t g_w (\bar{Q}\sigma^{\mu\nu} \tau^I t) \tilde{\varphi} W_{\mu\nu}^I$$

$$O_{tB} = y_t g_Y (\bar{Q}\sigma^{\mu\nu} t) \tilde{\varphi} B_{\mu\nu}$$

$$O_{tG} = y_t g_s (\bar{Q}\sigma^{\mu\nu} T^A t) \tilde{\varphi} G_{\mu\nu}^A,$$


$$O_{t\phi} = y_t^3 \left(\phi^\dagger \phi \right) (\bar{Q}t) \tilde{\phi}$$

see for example: Aguilar-Saavedra (arXiv:0811.3842)

Zhang and Willenbrock (arXiv:1008.3869)

+four-fermion operators

+non-top operators (mixing)



Top-quark operators & how to look for them

$$O_{\varphi Q}^{(3)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu^I \varphi \right) (\bar{Q}\gamma^\mu \tau^I Q)$$

$$O_{\varphi Q}^{(1)} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{Q}\gamma^\mu Q)$$

$$O_{\varphi t} = i\frac{1}{2}y_t^2 \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{t}\gamma^\mu t)$$

$$O_{tW} = y_t g_w (\bar{Q}\sigma^{\mu\nu} \tau^I t) \tilde{\varphi} W_{\mu\nu}^I$$

$$O_{tB} = y_t g_Y (\bar{Q}\sigma^{\mu\nu} t) \tilde{\varphi} B_{\mu\nu}$$

$$O_{tG} = y_t g_s (\bar{Q}\sigma^{\mu\nu} T^A t) \tilde{\varphi} G_{\mu\nu}^A,$$


$$O_{t\phi} = y_t^3 \left(\phi^\dagger \phi \right) (\bar{Q}t) \tilde{\phi}$$

see for example: Aguilar-Saavedra (arXiv:0811.3842)

Zhang and Willenbrock (arXiv:1008.3869)

+four-fermion operators

+non-top operators (mixing)




Operators entering various processes: Global approach needed

What's next?

Use SMEFT to
parametrise and look for
deviations from SM
predictions

What's next?


Use SMEFT to
parametrise and look for
deviations from SM
predictions



Use as many experimental
measurements as possible
Cross-sections+differential
distributions

What's next?

Use SMEFT to
parametrise and look for
deviations from SM
predictions




Use as many experimental
measurements as possible
Cross-sections+differential
distributions



Use the best SM
predictions
QCD/EW corrections

What's next?

Use SMEFT to
parametrise and look for
deviations from SM
predictions



Use as many experimental
measurements as possible
Cross-sections+differential
distributions




Use the best SM
predictions
QCD/EW corrections



Need for
precision also in
SMEFT

What's next?

Use SMEFT to
parametrise and look for
deviations from SM
predictions



Use as many experimental
measurements as possible
Cross-sections+differential
distributions



Use the best SM
predictions
QCD/EW corrections



Need for
precision also in
SMEFT



Need for precision
calculations
Automated tools
for the EFT



How can we improve EFT predictions?

- SMEFT@NLO
 - Mixing between operators: anomalous dimension matrix: [Jenkins et al arXiv:1308.2627](#), [1310.4838](#), [Alonso et al. 1312.2014](#)
 - Additional operators at NLO: e.g. chromomagnetic dipole in single top production

Recent progress in top physics:

- top pair [Franzosi and Zhang \(arxiv:1503.08841\)](#)
- single top [Zhang \(arxiv:1601.06163\)](#), [de Beurs, Laenen, Vreeswijk, EV \(arXiv:1807.03576\)](#)
- ttZ/γ [Bylund, Maltoni, Tsirikos, EV, Zhang \(arXiv:1601.08193\)](#), [Schulze and Rontsch \(arXiv:1404.1005\)](#)
- ttH [Maltoni, EV, Zhang \(arXiv:1607.05330\)](#)
- tZ/Hj [Degrande, Maltoni, Mimasu, EV, Zhang \(arXiv:1804.07773\)](#)

Single top production and decay




- The same EFT couplings enter both the production and decay
- The width of the top enters in the total cross-section calculation
- Computing the width of the top consistently is needed to match the Wbj and tj cross-sections

de Beurs, Laenen, Vreeswijk, EV arXiv:1807.03576


Single top production and decay

Beyond the narrow width approximation:

- Wbj production with off-shell and interference effects




- Resonant-aware matching to the Parton Shower ([arXiv: 1603.01178](https://arxiv.org/abs/1603.01178))
- W decay in MadSpin to retain spin information




top polarisation angles

$$\frac{1}{\sigma} \frac{d\sigma}{d \cos \theta_i^z} = \frac{1}{2} (1 + a_i P \cos \theta_i^z)$$

tZj/tHj associated production



Gauge-Higgs
Top couplings
TGC



\mathcal{O}_W	$\epsilon_{IJK} W_{\mu\nu}^I W^{J,\nu\rho} W_{\rho\mu}^K$	$\mathcal{O}_{\varphi Q}^{(3)}$	$i(\varphi^\dagger \overleftrightarrow{D}_\mu \tau_I \varphi)(\bar{Q} \gamma^\mu \tau^I Q) + \text{h.c.}$
$\mathcal{O}_{\varphi W}$	$(\varphi^\dagger \varphi - \frac{v^2}{2}) W_I^{\mu\nu} W_{\mu\nu}^I$	$\mathcal{O}_{\varphi Q}^{(1)}$	$i(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi)(\bar{Q} \gamma^\mu Q) + \text{h.c.}$
$\mathcal{O}_{\varphi WB}$	$(\varphi^\dagger \tau_I \varphi) B^{\mu\nu} W_{\mu\nu}^I$	$\mathcal{O}_{\varphi t}$	$i(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi)(\bar{t} \gamma^\mu t) + \text{h.c.}$
$\mathcal{O}_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^\dagger (\varphi^\dagger D_\mu \varphi)$	$\mathcal{O}_{\varphi tb}$	$i(\bar{\varphi} D_\mu \varphi)(\bar{t} \gamma^\mu b) + \text{h.c.}$
$\mathcal{O}_{\varphi \square}$	$(\varphi^\dagger \varphi) \square (\varphi^\dagger \varphi)$	$\mathcal{O}_{\varphi q}^{(1)}$	$i(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi)(\bar{q}_i \gamma^\mu q_i) + \text{h.c.}$
$\mathcal{O}_{t\varphi}$	$(\varphi^\dagger \varphi - \frac{v^2}{2}) \bar{Q} t \bar{\varphi} + \text{h.c.}$	$\mathcal{O}_{\varphi q}^{(3)}$	$i(\varphi^\dagger \overleftrightarrow{D}_\mu \tau_I \varphi)(\bar{q}_i \gamma^\mu \tau^I q_i) + \text{h.c.}$
\mathcal{O}_{tW}	$i(\bar{Q} \sigma^{\mu\nu} \tau_I t) \bar{\varphi} W_{\mu\nu}^I + \text{h.c.}$	$\mathcal{O}_{\varphi u}$	$i(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi)(\bar{u}_i \gamma^\mu u_i) + \text{h.c.}$
\mathcal{O}_{tB}	$i(\bar{Q} \sigma^{\mu\nu} t) \bar{\varphi} B_{\mu\nu} + \text{h.c.}$	$\mathcal{O}_{Qq}^{(3,1)}$	$(\bar{q}_i \gamma_\mu \tau_I q_i)(\bar{Q} \gamma^\mu \tau^I Q)$
\mathcal{O}_{tG}	$i(\bar{Q} \sigma^{\mu\nu} T_A t) \bar{\varphi} G_{\mu\nu}^A + \text{h.c.}$	$\mathcal{O}_{Qq}^{(3,8)}$	$(\bar{q}_i \gamma_\mu \tau_I T_A q_i)(\bar{Q} \gamma^\mu \tau^I T^A Q)$

Unique interplay

Pure gauge operators (4): $\mathcal{O}_{\varphi W}, \mathcal{O}_W, \mathcal{O}_{HW}, \mathcal{O}_{HB},$


Two-fermion top-quark operators (8): $\mathcal{O}_{\varphi Q}^{(3)}, \mathcal{O}_{\varphi Q}^{(1)}, \mathcal{O}_{\varphi t}, \mathcal{O}_{tW}, \mathcal{O}_{tB}, \mathcal{O}_{tG}, \mathcal{O}_{\varphi tb}, \mathcal{O}_{t\varphi}$

Four-fermion top-quark operators (2): $\mathcal{O}_{Qq}^{(3,1)}, \mathcal{O}_{Qq}^{(3,8)}.$

Helicity amplitudes for subprocesses

bW → tH

$\lambda_b, \lambda_W, \lambda_t$	SM	$\mathcal{O}_{t\varphi}$	$\mathcal{O}_{\varphi Q}^{(3)}$	$\mathcal{O}_{\varphi W}$	\mathcal{O}_{tW}	\mathcal{O}_{HW}
- , 0 , -	s^0	s^0	$\sqrt{s(s+t)}$	s^0	s^0	$\sqrt{s(s+t)}$
- , 0 , +	$\frac{1}{\sqrt{s}}$	$m_t\sqrt{-t}$	$m_t\sqrt{-t}$	$\frac{1}{\sqrt{s}}$	$\frac{m_W s}{\sqrt{-t}}$	$\frac{1}{\sqrt{s}}$
- , - , -	$\frac{1}{\sqrt{s}}$	$\frac{1}{\sqrt{s}}$	$m_W\sqrt{-t}$	$\frac{m_W s}{\sqrt{-t}}$	$m_t\sqrt{-t}$	$\frac{m_W(s+t)}{\sqrt{-t}}$
- , - , +	$\frac{1}{s}$	s^0	s^0	-	$\sqrt{s(s+t)}$	$\frac{1}{s}$
- , + , -	$\frac{1}{\sqrt{s}}$	-	$\frac{1}{\sqrt{s}}$	$\frac{m_W(s+t)}{\sqrt{-t}}$	$\frac{1}{\sqrt{s}}$	$\frac{m_W(s+t)}{\sqrt{-t}}$
- , + , +	s^0	-	s^0	s^0	s^0	$\frac{1}{s}$



bW → tZ



Amplitudes growing with energy as SM cancellations get spoiled →

Large deviations
Differential distributions

$\lambda_b, \lambda_W, \lambda_t, \lambda_Z$	SM	$\mathcal{O}_{\varphi Q}^{(3)}$	$\mathcal{O}_{\varphi Q}^{(1)}$	$\mathcal{O}_{\varphi t}$	\mathcal{O}_{tB}	\mathcal{O}_{tW}	\mathcal{O}_W	\mathcal{O}_{HW}	\mathcal{O}_{HB}
- , 0 , - , 0	s^0	$\sqrt{s(s+t)}$	-	-	-	s^0	s^0	$\sqrt{s(s+t)}$	s^0
- , 0 , + , 0	$\frac{1}{\sqrt{s}}$	$m_t\sqrt{-t}$	$m_t\sqrt{-t}$	$m_t\sqrt{-t}$	$m_Z\sqrt{-t}$	$\frac{m_W(2s+3t)}{\sqrt{-t}}$	-	$m_t\sqrt{-t}$	$m_t\sqrt{-t}$
- , - , - , 0	$\frac{1}{\sqrt{s}}$	$m_W\sqrt{-t}$	-	-	-	-	$\frac{m_W(s+2t)}{\sqrt{-t}}$	$m_W\sqrt{-t}$	$\frac{1}{\sqrt{s}}$
- , - , + , 0	$\frac{1}{s}$	s^0	s^0	s^0	s^0	$\sqrt{s(s+t)}$	s^0	s^0	$\frac{1}{\sqrt{s}}$
- , 0 , - , -	$\frac{1}{\sqrt{s}}$	$m_W\sqrt{-t}$	-	-	$m_t\sqrt{-t}$	$m_t\sqrt{-t}$	$\frac{m_W(s+2t)}{\sqrt{-t}}$	$\frac{m_W(ss_W^2+2t)}{\sqrt{-t}}$	$\frac{m_W s}{\sqrt{-t}}$
- , 0 , - , +	$\frac{1}{\sqrt{s}}$	-	-	-	-	-	$\frac{m_W(s+t)}{\sqrt{-t}}$	$\frac{m_W(s+t)}{\sqrt{-t}}$	$\frac{m_W(s+t)}{\sqrt{-t}}$
- , 0 , + , -	s^0	s^0	s^0	-	-	s^0	s^0	s^0	s^0
- , 0 , + , +	$\frac{1}{s}$	s^0	s^0	s^0	$\sqrt{s(s+t)}$	$\sqrt{s(s+t)}$	-	s^0	s^0
- , + , - , 0	$\frac{1}{\sqrt{s}}$	-	-	-	-	-	$\frac{m_W(s+t)}{\sqrt{-t}}$	$\frac{1}{\sqrt{s}}$	$\frac{1}{\sqrt{s}}$
- , + , + , 0	s^0	s^0	-	-	-	s^0	-	s^0	$\frac{1}{s}$
- , - , - , -	s^0	s^0	s^0	-	s^0	s^0	s^0	s^0	s^0
- , - , - , +	$\frac{1}{s}$	-	-	-	-	-	$\sqrt{s(s+t)}$	s^0	s^0
- , - , + , -	$\frac{1}{\sqrt{s}}$	-	-	-	-	$\frac{m_Z(s_W^2 t - 3c_W^2(2s+t))}{\sqrt{-t}}$	-	$\frac{1}{\sqrt{s}}$	$\frac{1}{\sqrt{s}}$
- , - , + , +	-	-	-	-	$m_W\sqrt{-t}$	$m_Z\sqrt{-t}$	$m_t\sqrt{-t}$	$m_t\sqrt{-t}$	$m_t\sqrt{-t}$
- , + , - , -	$\frac{1}{s}$	-	-	-	-	-	$\sqrt{s(s+t)}$	s^0	s^0
- , + , - , +	s^0	s^0	s^0	-	-	-	-	s^0	s^0
- , + , + , -	$\frac{1}{\sqrt{s}}$	-	-	-	-	-	$m_t\sqrt{-t}$	$m_t\sqrt{-t}$	$m_t\sqrt{-t}$
- , + , + , +	$\frac{1}{\sqrt{s}}$	-	-	-	-	$\frac{m_W(s+t)}{\sqrt{-t}}$	-	$\frac{1}{\sqrt{s}}$	$\frac{1}{\sqrt{s}}$

Differential results

tZj




Degrande, Maltoni, Mimasu, EV, Zhang arXiv:1804.07773

Large deviations in the tails, as expected from helicity amplitudes

Comparison with single top


	tj	tj ($p_T^t > 350$ GeV)	tZj	tZj ($p_T^t > 250$ GeV)
σ_{SM}	224 pb	880 fb	839 fb	69 fb
r_{tW}	0.0275	0.024	0.016	0.010
$r_{tW,tW}$	0.0162	0.35	0.095	0.67
$r_{\varphi Q^{(3)}}$	0.121	0.121	0.192	0.172
$r_{\varphi Q^{(3)},\varphi Q^{(3)}}$	0.0037	0.0037	0.029	0.114
$r_{\varphi tb,\varphi tb}$	0.00090	0.0008	0.0050	0.027
$r_{Qq^{(3,1)}}$	-0.353	-4.4	-0.59	-2.22
$r_{Qq^{(3,1)},Qq^{(3,1)}}$	0.126	11.5	0.65	5.1
$r_{Qq^{(3,8)},Qq^{(3,8)}}$	0.0308	2.73	0.133	1.01




- Increased sensitivity for dipoles and right-handed current (as expected from helicity analysis)
- 4-fermion operators sensitivity due to higher thresholds can be outperformed by high- p_T single top measurements

Current and future sensitivity

Current:



Future:



Degrande, Maltoni, Mimasu, EV, Zhang arXiv:1804.07773

tZj measurements:

CMS; PLB 779 (2018) 358-384: 0.75 ± 0.27

ATLAS; CERN-EP-2017-188: 1.31 ± 0.47

Promising for weak dipoles, RHCC and SU(2) current in particular for HL-LHC where high pT data can be used

Rare processes can play a role in a global fit

Towards a complete implementation@NLO

Aim to obtain a complete Monte Carlo implementation based on:

- Warsaw basis
- Degrees of freedom for top operators as in arXiv:1802.07237 (LHCTopWG)

Current status:

- 73 degrees of freedom (top, Higgs, gauge):
 - CP-conserving
 - Flavour assumption: $U(2)_Q \times U(2)_u \times U(3)_d \times U(3)_L \times U(3)_e$
- 0/2F@NLO operators validated (with previous partial NLO implementations)  <http://feynrules.irmp.ucl.ac.be/wiki/SMEFTatNLO>
- 4F@NLO operators validation: on-going

Future plans

- Full NLO model release (4F@NLO)
- Other flavour assumptions
- CP-violating effects

Work in progress with: C. Degrande, G. Durieux, F. Maltoni, K. Mimasu, C. Zhang

In practice


UFO model with UV+R2 counterterms

Import to MG5_aMC@NLO

Proceed as in SM case

```
MG5_aMC>import model EFT
MG5_aMC>generate p p > t t~ z NP=2 [QCD]
MG5_aMC>output
MG5_aMC>launch
```

In practice



Behind the scenes...

In practice

UFO model with UV+R2 counterterms
Import to MG5_aMC@NLO
Proceed as in SM case

```
MG5_aMC>import model EFT
MG5_aMC>generate p p > t t~ z NP=2 [QCD]
MG5_aMC>output
MG5_aMC>launch
```

Results:

Fixed order NLO

NLO+PS with MC@NLO

Implementation gives:


$$\sigma = \sigma_{\text{SM}} + \sum_i \frac{1\text{TeV}^2}{\Lambda^2} C_i \sigma_i + \sum_{i \leq j} \frac{1\text{TeV}^4}{\Lambda^4} C_i C_j \sigma_{ij}.$$

interference
with SM

interference between
operators, squared
contributions

A first application: A global top fit@NLO

Class	Notation	Degree of Freedom	Operator Definition
QQQQ	OQQ1	c_{QQ}^1	$2C_{qq}^{1(3333)} - \frac{2}{3}C_{qq}^{3(3333)}$
	OQQ8	c_{QQ}^8	$8C_{qq}^{3(3333)}$
	OQt1	c_{Qt}^1	$C_{qu}^{1(3333)}$
	OQt8	c_{Qt}^8	$C_{qu}^{8(3333)}$
	OQb1	c_{Qd}^1	$C_{qd}^{1(3333)}$
	OQb8	c_{Qd}^8	$C_{qd}^{8(3333)}$
	Ott1	c_{tt}^1	$C_{uu}^{1(3333)}$
	Otb1	c_{tb}^1	$C_{ud}^{1(3333)}$
	Otb8	c_{tb}^8	$C_{ud}^{8(3333)}$
	OQtQb1	c_{QtQb}^1	$C_{quqd}^{1(3333)}$
OQtQb8	c_{QtQb}^8	$C_{quqd}^{8(3333)}$	
QQqq	O81qq	$c_{Qq}^{1,8}$	$C_{qq}^{1(1331)} + 3C_{qq}^{3(1331)}$
	O11qq	$c_{Qq}^{1,1}$	$C_{qq}^{1(1331)} + \frac{1}{6}C_{qq}^{1(1331)} + \frac{1}{2}C_{qq}^{3(1331)}$
	O83qq	$c_{Qq}^{3,8}$	$C_{qq}^{1(1331)} - C_{qq}^{3(1331)}$
	O13qq	$c_{Qq}^{3,1}$	$C_{qq}^{3(1331)} + \frac{1}{6}(C_{qq}^{1(1331)} - C_{qq}^{3(1331)})$
	O8qt	c_{tq}^8	$C_{qu}^{8(1331)}$
	O1qt	c_{tq}^1	$C_{qu}^{1(1331)}$
	O8ut	c_{tu}^8	$2C_{uu}^{1(1331)}$
	O1ut	c_{tu}^1	$C_{uu}^{1(1331)} + \frac{1}{3}C_{uu}^{3(1331)}$
	O8qu	c_{Qu}^8	$C_{qu}^{8(3311)}$
	O1qu	c_{Qu}^1	$C_{qu}^{1(3311)}$
	O8dt	c_{td}^8	$C_{ud}^{8(3311)}$
	O1dt	c_{td}^1	$C_{ud}^{1(3311)}$
	O8qd	c_{Qd}^8	$C_{qd}^{8(3311)}$
	O1qd	c_{Qd}^1	$C_{qd}^{1(3311)}$
QQ + V, G, φ	OtG	c_{tG}	$\text{Re}\{C_{uG}^{(33)}\}$
	OtW	c_{tW}	$\text{Re}\{C_{uW}^{(33)}\}$
	OtB	c_{tB}	$\text{Re}\{C_{dW}^{(33)}\}$
	OtZ	c_{tZ}	$\text{Re}\{-s_W C_{uB}^{(33)} + c_W C_{uW}^{(33)}\}$
	Otf	$c_{\varphi tb}$	$\text{Re}\{C_{\varphi ud}^{(33)}\}$
	Ofq3	$c_{\varphi q}^3$	$C_{\varphi q}^{3(33)}$
	Opm	$c_{\varphi q}$	$C_{\varphi q}^{1(33)} - C_{\varphi q}^{3(33)}$
	Opt	$c_{\varphi t}$	$C_{\varphi u}^{(33)}$
	Otp	$c_{t\varphi}$	$\text{Re}\{C_{u\varphi}^{(33)}\}$




Rich phenomenology

34 d.o.f.
CP-conserving


Hartland, Maltoni, Nocera, Rojo, Slade, EV and Zhang, arXiv:1901.05965

The impact of multiple measurements



$$O_{Qq}^{1,8} = (\bar{Q}\gamma_\mu T^A Q)(\bar{q}_i\gamma^\mu T^A q_i)$$

$$O_{Qq}^{3,8} = (\bar{Q}\gamma_\mu T^A \tau^I Q)(\bar{q}_i\gamma^\mu T^A \tau^I q_i)$$



$$O_{tq}^8 = (\bar{q}_i\gamma^\mu T^A q_i)(\bar{t}\gamma_\mu T^A t)$$

$$O_{Qq}^{1,8} = (\bar{Q}\gamma_\mu T^A Q)(\bar{q}_i\gamma^\mu T^A q_i)$$

Brivio, Bruggisser, Maltoni, Moutafis, Plehn, EV, Westhoff, Zhang arXiv:1910.03606

Some considerations for a fit


- Validity of the EFT expansion: $E < \Lambda$
 - Ensure results are not dominated by high energy regions
 - report limits as a function of the max scale probed [Contino et al arXiv:1604.06444](#)
- Range of Wilson coefficients:
 - The theory: perturbativity, unitarity, linear or non-linear EFT, UV completion
 - The experimental limits: Think about and use as many processes as possible to extract allowed range
- $1/\Lambda^2$ vs $1/\Lambda^4$ contributions
 - $1/\Lambda^2$ suppressed due to helicity [Azatov et al arXiv:1607.05236](#)
 - $1/\Lambda^4$ can be large for loosely constrained operator coefficients/strongly coupled theories

$$C_i^2 \frac{E^4}{\Lambda^4} > C_i \frac{E^2}{\Lambda^2} > 1 > \frac{E^2}{\Lambda^2}$$

$E < \Lambda$ satisfied but $O(1/\Lambda^4)$ large for large operator coefficients

1/Λ² vs 1/Λ⁴ contributions some examples

1)




1/Λ² is not positive definite
1/Λ² is not suppressed PS point by PS point
1/Λ² is suppressed only when integrating over the PS

3) ttZ production

13TeV	\mathcal{O}_{tW}
$\sigma_{i,NLO}^{(1)}$	$-1.7(2)^{+31.3\%}_{-49.1\%}$
$\sigma_{ii,NLO}^{(2)}$	$24.2^{+8.2\%}_{-11.2\%}$


2)



1/Λ² is suppressed compared to 1/Λ⁴
1/Λ⁴ from dimension-6 much larger than interference of SM with dim-8

Reasons why the interference is suppressed:
1) An accidental cancellation between the contributions of the gg and qq channels
2) \mathcal{O}_{tW} not producing a longitudinal Z

Impact of quadratic terms in top production




$$O_{Qq}^{1,8} = (\bar{Q}\gamma_\mu T^A Q)(\bar{q}_i\gamma^\mu T^A q_i)$$

$$O_{Qq}^{3,8} = (\bar{Q}\gamma_\mu T^A \tau^I Q)(\bar{q}_i\gamma^\mu T^A \tau^I q_i)$$

Brivio, Bruggisser, Maltoni, Moutafis, Plehn, EV, Westhoff, Zhang arXiv:1910.03606

Impact of quadratic terms in top production



$$O_{tq}^8 = (\bar{q}_i \gamma^\mu T^A q_i) (\bar{t} \gamma_\mu T^A t)$$

$$O_{Qq}^{1,8} = (\bar{Q} \gamma_\mu T^A Q) (\bar{q}_i \gamma^\mu T^A q_i)$$

Brivio, Bruggisser, Maltoni, Moutafis, Plehn, EV, Westhoff, Zhang arXiv:1910.03606

Global fit Setup

Theory

(N)NLO QCD for SM
NLO QCD for SMEFT
State-of-the-art PDFs without top data

Data

Top pair production and single top (differential)
Associated production with W,Z,H
W helicity fractions
Parton-level

Global SMEFT fit
of the top-quark sector

Based on NNPDF
Faithful uncertainty estimate
Avoid under- and over-fitting
Validated on pseudo-data (closure test)

Methodology

Fit results can be used to bound
specific UV complete models
New data can be straightforwardly added
Plan to extend to Higgs, gauge sector etc

Output

Observables and theory predictions

Data

Dataset	n_{dat}
ATLAS_tt_8TeV_ljets [$m_{t\bar{t}}$]	7
CMS_tt_8TeV_ljets [y_t]	10
CMS_tt2D_8TeV_dilep [($m_{t\bar{t}}, y_t$)]	16
CMS_tt_13TeV_ljets2 [$y_{t\bar{t}}$]	8
CMS_tt_13TeV_dilep [$y_{t\bar{t}}$]	6
CMS_tt_13TeV_ljets_2016 [y_t]	11
ATLAS_WhelF_8TeV	3
CMS_WhelF_8TeV	3
<hr/>	
CMS_ttbb_13TeV	1
CMS_tttt_13TeV	1
ATLAS_tth_13TeV	1
CMS_tth_13TeV	1
ATLAS_ttZ_8TeV	1
ATLAS_ttZ_13TeV	1
CMS_ttZ_8TeV	1
CMS_ttZ_13TeV	1
ATLAS_ttW_8TeV	1
ATLAS_ttW_13TeV	1
CMS_ttW_8TeV	1
CMS_ttW_13TeV	1
<hr/>	
CMS_t_tch_8TeV_dif	6
ATLAS_t_tch_8TeV [y_t]	4
ATLAS_t_tch_8TeV [y_t]	4
ATLAS_t_sch_8TeV	1
CMS_t_tch_13TeV_dif [y_t]	4
CMS_t_sch_8TeV	1
ATLAS_tW_inc_8TeV	1
CMS_tW_inc_8TeV	1
ATLAS_tW_inc_13TeV	1
CMS_tW_inc_13TeV	1
ATLAS_tZ_inc_13TeV	1
CMS_tZ_inc_13TeV	1
<hr/>	
Total	102

Top-pair production
W-helicities

4 tops, ttbb, top-pair associated production

Single top
t-channel, s-channel, tW, tZ

One distribution from each dataset, to avoid double counting


Theoretical predictions

Process	SM	SMEFT
$t\bar{t}$	NNLO QCD	NLO QCD
single-t (t-ch)	NNLO QCD	NLO QCD
single-t (s-ch)	NLO QCD	NLO QCD
tW	NLO QCD	NLO QCD
tZ	NLO QCD	LO QCD + NLO SM K -factors
$t\bar{t}W(Z)$	NLO QCD	LO QCD + NLO SM K -factors
$t\bar{t}h$	NLO QCD	LO QCD + NLO SM K -factors
$t\bar{t}\bar{t}$	NLO QCD	LO QCD + NLO SM K -factors
$t\bar{t}b\bar{b}$	NLO QCD	LO QCD + NLO SM K -factors

Baseline fit includes:

- Best available SM predictions
- NLO EFT predictions
- $O(1/\Lambda^4)$ terms

Global top EFT fit@NLO




First limits reported for some operators

Improvement for some operators: e.g. O_{tG} , O_{83qq} , O_{bW}

Hartland, Maltoni, Nocera, Rojo, Slade, EV and Zhang, arXiv:1901.05965

Impact of higher-order terms

Fit allows to check the impact of NLO QCD corrections and of including the $O(1/\Lambda^4)$ terms



Non-trivial impact of the two effects, can be different operator-by-operator

Hartland, Maltoni, Nocera, Rojo, Slade, EV and Zhang, arXiv:1901.05965

Outline

- Introduction to the EFT
- EFT in top quark physics
 - Precision calculations in the EFT
 - Towards global fits in the top sector
- EFT in the top-Higgs sector
 - Top loops in the EFT

The top-Higgs interface

$$O_{t\phi} = y_t^3 (\phi^\dagger \phi) (\bar{Q}t) \tilde{\phi},$$

$$O_{\phi G} = y_t^2 (\phi^\dagger \phi) G_{\mu\nu}^A G^{A\mu\nu},$$


$$O_{tG} = y_t g_s (\bar{Q} \sigma^{\mu\nu} T^A t) \tilde{\phi} G_{\mu\nu}^A$$

See also

Degrande et al. arXiv:1205.1065

Grojean et al. arXiv:1312.3317

Azatov et al arXiv:1608.00977



ttH




H, H+j, HH

Use with 1) ttH and 2) H, H+j to break degeneracy between operators and extract maximal information on these operators




Maltoni, EV, Zhang: arXiv:1607.05330

SMEFT in single Higgs production

LO:




NLO:



13 TeV	σ LO	σ/σ_{SM} LO	σ NLO	σ/σ_{SM} NLO	K
σ_{SM}	$21.3^{+34.0+1.5\%}_{-25.0-1.5\%}$	1.0	$36.6^{+26.4+1.9\%}_{-20.0-1.6\%}$	1.0	1.71
σ_1	$-2.93^{+34.0+1.5\%}_{-25.0-1.5\%}$	-0.138	$-4.70^{+24.8+1.9\%}_{-20.0-1.6\%}$	-0.127	1.61
σ_2	$2660^{+34.0+1.5\%}_{-25.0-1.5\%}$	125	$4130^{+23.9+1.9\%}_{-19.6-1.6\%}$	114	1.55
σ_3	$50.5^{+34.0+1.5\%}_{-25.0-1.5\%}$	2.38	$83.5^{+26.0+1.9\%}_{-20.6-1.6\%}$	2.28	1.65


Deutschmann, Duhr, Maltoni, EV
arXiv:1708.00460

SMEFT in Higgs production



Higgs p_T


Deutschmann, Duhr, Maltoni, EV arXiv:1708.00460



Higgs p_T


Grazzini et al 1612.00283

SMEFT in Higgs production



Higgs p_T

Deutschmann, Duhr, Maltoni, EV arXiv:1708.00460




Higgs p_T

Grazzini et al 1612.00283

Non-trivial EFT parameter combinations not changing the inclusive cross-section but changing the distributions

Constraints using two-operator fits



Current limits using LHC measurements

$$O_{t\phi} = y_t^3 (\phi^\dagger \phi) (\bar{Q}t) \tilde{\phi}$$

$$O_{\phi G} = y_t^2 (\phi^\dagger \phi) G_{\mu\nu}^A G^{A\mu\nu}$$

14TeV projection
3000 fb⁻¹



Maltoni, EV, Zhang arXiv:
1607.05330

Double Higgs production

The Higgs potential

$$V(H) = \frac{1}{2}M_H^2 H^2 + \lambda_{HHH} v H^3 + \frac{1}{4}\lambda_{HHHH} H^4$$



$$\lambda_{HHH} = \lambda_{HHHH} = \frac{M_H^2}{2v^2}$$



Phys.Lett. B732 (2014) 142-149

A challenging process at the LHC

HH in the EFT



top Yukawa, ggh(h) coupling, top-gluon interaction, Higgs self-coupling


The present

Given the current constraints on $\sigma(\text{HH})$, $\sigma(\text{H})$ and the fresh ttH measurement, the Higgs self-coupling can be currently constrained “ignoring” other couplings

The future

Precise knowledge of other Wilson coefficients will be needed to bound λ as the bound gets closer to SM
Differential distributions will also be necessary

HH in the EFT



The present

Given the current constraints on $\sigma(HH)$, $\sigma(H)$ and the fresh ttH measurement, the Higgs self-coupling can be currently constrained “ignoring” other couplings

$$O_{t\phi} = y_t^3 (\phi^\dagger \phi) (\bar{Q}t) \tilde{\phi},$$

$$O_{\phi G} = y_t^2 (\phi^\dagger \phi) G_{\mu\nu}^A G^{A\mu\nu},$$

$$O_{tG} = y_t g_s (\bar{Q} \sigma^{\mu\nu} T^A t) \tilde{\phi} G_{\mu\nu}^A$$

$$O_6 = -\lambda (\phi^\dagger \phi)^3$$

$$O_H = \frac{1}{2} (\partial_\mu (\phi^\dagger \phi))^2$$


top Yukawa, $ggh(h)$ coupling, top-gluon interaction, Higgs self-coupling

The future

Precise knowledge of other Wilson coefficients will be needed to bound λ as the bound gets closer to SM

Differential distributions will also be necessary

HH in the EFT



$$O_{t\phi} = y_t^3 (\phi^\dagger \phi) (\bar{Q}t) \tilde{\phi},$$

$$O_{\phi G} = y_t^2 (\phi^\dagger \phi) G_{\mu\nu}^A G^{A\mu\nu},$$

$$O_{tG} = y_t g_s (\bar{Q} \sigma^{\mu\nu} T^A t) \tilde{\phi} G_{\mu\nu}^A$$

$$O_6 = -\lambda (\phi^\dagger \phi)^3$$

$$O_H = \frac{1}{2} (\partial_\mu (\phi^\dagger \phi))^2$$

top Yukawa, ggh(h) coupling, top-gluon interaction, Higgs self-coupling

The present


Given the current constraints on $\sigma(\text{HH})$, $\sigma(\text{H})$ and the fresh $t\bar{t}H$ measurement, the Higgs self-coupling can be currently constrained “ignoring” other couplings

The future


Precise knowledge of other Wilson coefficients will be needed to bound λ as the bound gets closer to SM
Differential distributions will also be necessary

Going beyond QCD corrections in the EFT

Are we measuring




or



?



NLO EW in SMEFT may not be small:

$$\mathcal{O}(\alpha_{EW}/\pi \cdot C_t/C_H) \quad \text{instead of} \quad \mathcal{O}(\alpha_{EW}/\pi)$$




Weak corrections can be important for unconstrained operators

Towards weak loops in the EFT



$$\begin{aligned}
 O_{t\varphi} &= \bar{Q}t\tilde{\varphi}(\varphi^\dagger\varphi) + h.c., \\
 O_{\varphi Q}^{(3)} &= (\varphi^\dagger i\overleftrightarrow{D}_\mu^I\varphi)(\bar{Q}\gamma^\mu\tau^I Q), \\
 O_{\varphi tb} &= (\tilde{\varphi}^\dagger iD_\mu\varphi)(\bar{t}\gamma^\mu b) + h.c., \\
 O_{tB} &= (\bar{Q}\sigma^{\mu\nu}t)\tilde{\varphi}B_{\mu\nu} + h.c., \\
 O_{\varphi t} &= (\varphi^\dagger i\overleftrightarrow{D}_\mu\varphi)(\bar{t}\gamma^\mu t), \\
 O_{\varphi Q}^{(1)} &= (\varphi^\dagger i\overleftrightarrow{D}_\mu\varphi)(\bar{Q}\gamma^\mu Q), \\
 O_{tW} &= (\bar{Q}\sigma^{\mu\nu}\tau^I t)\tilde{\varphi}W_{\mu\nu}^I + h.c.,
 \end{aligned}$$

Current constraints from top LHC measurements



Poor knowledge of top couplings leads to uncertainties on Higgs measurements at the LHC:

	$\gamma\gamma$	γZ	bb	WW*	ZZ*
gg	(-100%, 1980%)	(-88%, 200%)	(-40%, 48%)	(-40%, 47%)	(-40%, 46%)
VBF	(-100%, 1880%)	(-88%, 170%)	(-6.1%, 5.3%)	(-6.8%, 6.7%)	(-8.8%, 9.2%)
WH	(-100%, 1880%)	(-88%, 170%)	(-5.5%, 4.2%)	(-6.1%, 5.6%)	(-7.8%, 7.9%)
ZH	(-100%, 1880%)	(-87%, 170%)	(-6.5%, 5.9%)	(-7.1%, 7.1%)	(-9.4%, 9.9%)


loop-induced

tree-level


EV, Zhang arXiv:1804.09766

Weak loops in the EFT: Future colliders

Circular Electron Positron Collider & HL-LHC: Top + Higgs Global Fit





Higgs production and decay



WW production

precision of the Higgs parameters at CC (global fit, $\Delta\chi^2=1$)



Durieux, Gu, EV, Zhang arXiv:1809.03520

Outlook

- SMEFT is a consistent way to look for new interactions
- Higher-order corrections needed to match SM precision and experimental accuracy
- Progress in top-quark processes: single top, $t(t)+V$, $t(t)+H$ as well as in the Monte Carlo automation of these corrections
- QCD corrections important both for total cross-sections and distributions: SM k-factors are not enough
- First global fits results already available: important to include NLO predictions where available and to combine as many processes as possible to extract maximal information
- Electroweak loop corrections can be important, progress towards computing them and assessing their impact

Thank you for your attention