8. Cross section

idea:



we repeat this experiment N times with randomly distributed impact parameter $\vec{\varrho}$ (area A larger than dimensions of the object)

N_{sc} = number of scattered particles

$$\Rightarrow \quad \frac{N_{sc}}{N} = \frac{\sigma}{A}$$

σ = cross section (in this case geometric cross section of object $\perp \vec{p}$)

$$N_{sc} = \underbrace{\frac{N}{A}}_{n} \sigma$$

n = number of incoming particles per area

more detailed information from observation of angular distribution of scattered particles.



$N_{sc}(\Delta\Omega)$ = number of particles scattered into solid angle $\Delta\Omega$

$$N_{sc}(\Delta\Omega) = n\,\sigma(\Delta\Omega)$$

infinitesimal: $\sigma(d\Omega) = \underbrace{\frac{d\sigma}{d\Omega}}_{\text{differential cross section}} d\Omega$

concept of cross section in particle physics:



mean values of (three-) momenta

$|\phi\rangle \qquad\qquad |\chi\rangle$

$$|\psi \text{ in}\rangle = |\phi\rangle \otimes |\chi\rangle$$

particles <u>distinguishable</u>

(identical particles $\rightarrow$ see later)

S matrix element $\langle k_1, \dots, k_n \text{ out} | \psi \text{ in}\rangle =$

$$= \int d\mu(p)\, d\mu(q)\, \underbrace{\langle k_1, \dots, k_n \text{ out}}_{n \text{ distinguishable particles}} | p, q \text{ in}\rangle \phi(p)\, \chi(q)$$

(identical particles $\rightarrow$ see later); spin indices suppressed

$$\langle k_1, \dots, k_n \text{ out} \,|\, p, q \text{ in} \rangle =$$

$$= i\,(2\pi)^4\, \delta^{(4)}\Big(p+q - \sum_{i=1}^{n} k_i\Big)\, \mathcal{M}(p,q \to k_1, \dots, k_n)$$

$$w\,(|\psi\rangle \to B) = \int_B d\mu(k_1) \dots d\mu(k_n)\, |\langle k_1, \dots, k_n \text{ out} | \psi \text{ in}\rangle|^2$$

region in n-particle "phase-space"

$$= \int d\mu(k_1) \dots d\mu(k_n)\, \langle \psi \text{ in} | k_1, \dots, k_n \text{ out}\rangle \langle k_1, \dots, k_n \text{ out} | \psi \text{ in}\rangle$$

$= \langle \psi \text{ in} | P_B | \psi \text{ in} \rangle =$ expectation value of P_B in the state $|\psi \text{ in}\rangle$

$$P_B := \int d\mu(k_1) \dots d\mu(k_n)\, |k_1, \dots, k_n \text{ out}\rangle \langle k_1, \dots, k_n \text{ out}|$$

projector on region B of n-particle phase-space ($\to$ generalization in the case of some identical particles in the final state now obvious)

$$N_{sc}(|\psi\rangle \to B) = \underbrace{\frac{N}{A}}_{n} \underbrace{\int_A d^2\xi \, w(|\phi_{\vec{\xi}}\rangle|\chi\rangle \to B)}_{=:\ \sigma(|\psi\rangle \to B)}$$

$$\phi_{\vec{\xi}}(p) = e^{-i\vec{p}\cdot\vec{\xi}}\,\phi(p)$$

$$\sigma(|\psi\rangle \to B) = \int d^2\xi \, w(|\phi_{\vec{\xi}}\rangle|\chi\rangle \to B)$$

$$= \int_{A\to\mathbb{R}^2} d^2\xi \int_B d\mu(k_1)\dots d\mu(k_n) \int d\mu(p)\,d\mu(q)\,\langle k_1,\dots,k_n\,\mathrm{out}|p,q\,\mathrm{in}\rangle$$

$$e^{-i\vec{p}\cdot\vec{\xi}}\,\phi(p)\,\chi(q) \int d\mu(p')\,d\mu(q')\,\langle k_1,\dots,k_n\,\mathrm{out}|p',q'\,\mathrm{in}\rangle^*$$

$$e^{+i\vec{p}'\cdot\vec{\xi}}\,\phi(p')^*\,\chi(q')^*$$

$$= \int_B d\mu(k_1)\dots d\mu(k_n) \int d\mu(p)\,d\mu(q)\,d\mu(p')\,d\mu(q')$$

$$(2\pi)^2\,\delta^{(2)}(\underbrace{\vec{p}_\perp - \vec{p}_\perp{}'}_{\perp\,\vec{p}_{1,2}})\,\phi(p)\,\phi(p')^*\,\chi(q)\,\chi(q')^*$$

$$(2\pi)^4\,\delta^{(4)}\Big(p+q-\sum_{i=1}^n k_i\Big)\,\mathcal{M}(p,q\to k_1,\dots,k_n)$$

$$(2\pi)^4\,\delta^{(4)}\Big(p'+q'-\sum_{i=1}^n k_i\Big)\,\mathcal{M}(p',q'\to k_1,\dots,k_n)^*$$

$$= (2\pi)^{10} \int d\mu(k_1) \dots d\mu(k_n) \int d\mu(p)\, d\mu(q)$$

$$\delta^{(4)}\left(p+q-\sum_{i=1}^{n} k_i\right) \underbrace{\int d\mu(p')\, d\mu(q')}_{\text{six integrations}}$$

$$\underbrace{\delta^{(4)}(p'+q'-p-q)\ \delta^{(2)}(\vec{p}_\perp - \vec{p}_\perp{}')}_{\text{six } \delta\text{-functions}}$$

$$\mathcal{M}(p,q \to k_1,\dots,k_n)\ \mathcal{M}(p',q' \to k_1,\dots,k_n)^*$$

$$\Phi(p)\ \Phi(p')^*\ \chi(q)\ \chi(q')^*$$

$\vec{p}_1 = |\vec{p}_1|\, \vec{e}_z$:

$$\delta^{(4)}(p'+q'-p-q)\ \delta^{(2)}(\vec{p}_\perp - \vec{p}_\perp{}')$$

$$= \delta\left(\sqrt{\vec{p}'^2+m_1^2} + \sqrt{\vec{q}'^2+m_2^2} - \sqrt{\vec{p}^2+m_1^2} - \sqrt{\vec{q}^2+m_2^2}\right)$$

$$\delta(p_x'+q_x'-p_x-q_x)\ \delta(p_y'+q_y'-p_y-q_y)$$

$$\delta(p_z'+q_z'-p_z-q_z)\ \delta(p_x-p_x')\ \delta(p_y-p_y')$$

$$= \delta^{(2)}(\vec{p}_\perp - \vec{p}_\perp{}') \; \delta^{(2)}(\vec{q}_\perp - \vec{q}_\perp{}')$$

$$\delta(p_z' + q_z' - p_z - q_z)$$

$$\delta\Big(\sqrt{p_z'^2 + \vec{p}_\perp^{\,2} + m_1^2} + \sqrt{q_z'^2 + \vec{q}_\perp^{\,2} + m_2^2} -$$

$$- \sqrt{p_z^2 + \vec{p}_\perp^{\,2} + m_1^2} - \sqrt{q_z^2 + \vec{q}_\perp^{\,2} + m_2^2}\Big) = (*)$$

$\rightarrow$ system of equations

$$p_z' + q_z' = p_z + q_z$$

$$\sqrt{p_z'^2 + M_1^2} + \sqrt{q_z'^2 + M_2^2} = \sqrt{p_z^2 + M_1^2} + \sqrt{q_z^2 + M_2^2}$$

where $M_1^2 = \vec{p}_\perp^{\,2} + m_1^2$, $M_2^2 = \vec{q}_\perp^{\,2} + m_2^2$

relevant solution: $p_z' = p_z$, $q_z' = q_z$

other solution possible, but contribution vanishes when multiplied with wave functions in momentum space (concentrated around $\vec{p}_1$ and $\vec{p}_2$, respectively)

we have an expression of the form

$$\delta[f(x,y)]\ \delta[g(x,y)]$$

where $f(a,b)=0, \quad g(a,b)=0$

$$\overset{\text{ex.}}{\Rightarrow} \quad \delta[f(x,y)]\ \delta[g(x,y)] =$$

$$= \frac{\delta(x-a)\ \delta(x-b)}{\left|\det\begin{pmatrix} f_{,x} & g_{,x} \\ f_{,y} & g_{,y}\end{pmatrix}_{\substack{x=a\\y=b}}\right|} + \dots$$

contributions related to further solutions of $f(x,y)=g(x,y)=0$

in our case, we obtain:

$$\delta(p_z'+q_z'-p_z-q_z)\ \delta\left(\sqrt{p_z'^2+M_1^2}+\sqrt{q_z'^2+M_2^2}-\sqrt{p_z^2+M_1^2}-\sqrt{q_z^2+M_2^2}\right)$$

$$= \frac{\delta(p_z'-p_z)\ \delta(q_z'-q_z)}{\left|\frac{q_z}{q^0}-\frac{p_z}{p^0}\right|} + \dots$$

$$\Rightarrow \quad (*) = \frac{\delta^{(3)}(\vec{p}\,' - \vec{p})\ \delta^{(3)}(\vec{q}\,' - \vec{q})}{\left| \frac{p_{\parallel}}{p^0} - \frac{q_{\parallel}}{q^0} \right|}$$

$\parallel$ means: $\parallel$ to $\vec{p}_1$

$$\Rightarrow \quad \sigma\left(|\psi\rangle \to B\right) = (2\pi)^{10} \int_B d\mu(k_1) \dots d\mu(k_n)$$

$$\int d\mu(p)\, d\mu(q)\ \delta^{(4)}\left(p + q - \sum_{i=1}^{n} k_i\right)$$

$$\int d\mu(p')\, d\mu(q')\ \frac{\delta^{(3)}(\vec{p}\,' - \vec{p})\ \delta^{(3)}(\vec{q}\,' - \vec{q})}{\left| \frac{p_{\parallel}}{p^0} - \frac{q_{\parallel}}{q^0} \right|}$$

$$\mathcal{M}(p, q \to k_1, \dots, k_n)\ \mathcal{M}(p', q' \to k_1, \dots, k_n)^*$$

$$\Phi(p)\ \Phi(p')^*\ \chi(q)\ \chi(q')^* =$$

$$= \int_B d\mu(k_1) \dots d\mu(k_n) \int d\mu(p)\, d\mu(q)\ (2\pi)^4 \delta^{(4)}\left(p + q - \sum_{i=1}^{n} k_i\right)$$

$$\frac{1}{2p^0}\ \frac{1}{2q^0}\ \frac{1}{\left| \frac{p_{\parallel}}{p^0} - \frac{q_{\parallel}}{q^0} \right|}\ \left|\mathcal{M}(p, q \to k_1, \dots, k_n)\right|^2\ |\Phi(p)|^2\, |\chi(q)|^2$$

<u>if</u> $|\phi(p)|^2$ sufficiently <u>concentrated</u> around $\vec{p}_1$

and $|\chi(q)|^2$ —"— —"— —"— $\vec{p}_2$

(relative to variations of $|\mathcal{M}(p,q \to k_1,\dots,k_n)|^2$)

$$\longrightarrow \quad \sigma(|\psi\rangle \to B) \simeq$$

$$\simeq \int_B d\mu(k_1)\dots d\mu(k_n)\,(2\pi)^4\,\delta^{(4)}\Big(p_1+p_2-\sum_{i=1}^{n} k_i\Big)$$

$$\frac{1}{4\,p_1^0\,p_2^0}\;\underbrace{\frac{1}{\dfrac{|\vec{p}_1|}{p_1^0}+\dfrac{|\vec{p}_2|}{p_2^0}}}_{|\vec{v}_1-\vec{v}_2|}\;|\mathcal{M}(p_1,p_2\to k_1,\dots,k_n)|^2$$

$$\underbrace{\int d\mu(p)\,|\phi(p)|^2}_{1}\;\underbrace{\int d\mu(q)\,|\chi(q)|^2}_{1}$$

<u>independent</u> of the exact form of ϕ and χ

$$\sigma(p_1,p_2\to B)=\int_B d\mu(k_1)\dots d\mu(k_n)\,(2\pi)^4\,\delta^{(4)}\Big(p_1+p_2-\sum_{i=1}^{n} k_i\Big)\,\frac{|\mathcal{M}|^2}{4\,(|\vec{p}_1|\,p_2^0+|\vec{p}_2|\,p_1^0)}$$

remark: $|\vec{p}_1|\,p_2^0+|\vec{p}_2|\,p_1^0=\sqrt{(p_1\cdot p_2)^2-m_1^2m_2^2}$ Lorentz invariant

<u>identical</u> particles in initial state

$\phi(p)$ one-particle (momentum space) wave function concentrated around $\vec{p} = \vec{p}_1$ $\left(\int d\mu(p)\, |\phi(p)|^2 = 1 \right)$

analogously: $\chi(q)$ one-particle wave function concentrated around $\vec{q} = \vec{p}_2$ $\left(\int d\mu(q)\, |\chi(q)|^2 = 1 \right)$

in momentum space:



$$\langle \phi | \chi \rangle = \int d\mu(p)\, \chi(p)^* \phi(p) = 0$$

(no overlap)

$$\Rightarrow |\psi\, in\rangle = \frac{1}{2}\int d\mu(p)\, d\mu(q)\, |p,q\, in\rangle\, [\phi(p)\,\chi(q) \pm \chi(p)\,\phi(q)]$$

(minus sign for fermions) is a properly normalized two-particle state

$$|p,q\, in\rangle = \pm\, |q,p\, in\rangle \Rightarrow |\psi\, in\rangle = \int d\mu(p)\, d\mu(q)\, |p,q\, in\rangle\, \phi(p)\,\chi(q)$$

→ further calculation <u>exactly</u> <u>the same</u> as for distinguishable particles

→ result was to be expected, as <u>no</u> overlap of momentum-space wave functions $\phi(p)$, $\chi(p)$ (particles can be distinguished <u>before</u> the interaction)

<u>identical</u> particles in <u>final state</u>:

consider first the case where <u>all</u> n final state particles identical

$$\rightarrow P_B = \frac{1}{n!} \int_B d\mu(k_1) \ldots d\mu(k_n)\, |k_1, \ldots, k_n\, out\rangle\langle k_1, \ldots, k_n\, out|$$

domain of integration B <u>symmetric</u> with respect to permutations of $k_1, \ldots, k_n$

<u>statistical factor</u> $\frac{1}{n!}$ in cross section formula

general case:

n_1 identical particles of type 1 (momenta $k_1^{(1)} \dots k_{n_1}^{(1)}$)

. . . . . .

n_ℓ —"— —"— —"— —"— ℓ (momenta $k_1^{(\ell)} \dots k_{n_\ell}^{(\ell)}$)

. . . . . .

n_r —"— —"— —"— —"— r (momenta $k_1^{(r)}, \dots, k_{n_r}^{(r)}$)

$$\sum_{\ell=1}^{r} n_\ell = n \quad \rightarrow \text{statistical factor } S = \prod_{\ell=1}^{r} \frac{1}{n_\ell!}$$

domain of integration B totally symmetric with respect to permutations of $k_1^{(\ell)}, \dots, k_{n_\ell}^{(\ell)}$ $(\ell = 1, \dots r)$

summary: $\langle k_1, \dots, k_n \text{ out} | p_1, p_2 \text{ in} \rangle =$

$$= i\,(2\pi)^4\, \delta^{(4)}\Big(p_1 + p_2 - \sum_{i=1}^{n} k_i\Big)\; \mathcal{M}(p_1, p_2 \rightarrow k_1, \dots, k_n)$$

$$\sigma(p_1, p_2 \rightarrow B) =$$

$$= \frac{S}{4\sqrt{(p_1 \cdot p_2)^2 - m_1^2 m_2^2}} \int_B d\mu(k_1) \dots d\mu(k_n)\, (2\pi)^4\, \delta^{(4)}\Big(p_1 + p_2 - \sum_{i=1}^{n} k_i\Big) \times |\mathcal{M}(p_1, p_2 \rightarrow k_1, \dots, k_n)|^2$$

application of cross section formula to scattering of identical scalars $\varphi(p_1)\varphi(p_2) \to \varphi(p_3)\varphi(p_4)$

$$(p_1+p_2)^2 = s \quad \Rightarrow \quad 2m^2 + 2p_1 p_2 = s$$

$$\Rightarrow \quad p_1 \cdot p_2 = \frac{s}{2} - m^2$$

$$\Rightarrow \quad (p_1 \cdot p_2)^2 - m^4 = \frac{s^2}{4} - s m^2 = \frac{s}{4}(s-4m^2)$$

$$\sigma = \frac{1}{2!} \frac{1}{4\sqrt{\frac{s}{4}(s-4m^2)}} \int d\mu(p_3)\, d\mu(p_4)\, (2\pi)^4 \delta^{(4)}(p_1+p_2-p_3-p_4) \times |\mathcal{M}(s,t)|^2$$

$$= \frac{1}{4\sqrt{s}\sqrt{s-4m^2}} \int \frac{d^3p_3}{(2\pi)^3 2p_3^0} \frac{d^3p_4}{(2\pi)^4 2p_4^0} (2\pi)^4 \delta^{(4)}(p_1+p_2-p_3-p_4) \times |\mathcal{M}(s,t)|^2$$

$$= \frac{1}{4(2\pi)^2\sqrt{s}\sqrt{s-4m^2}} \int \frac{d^3p_3}{2p_3^0}\, d^4p_4\, \Theta(p_4^0)\, \delta(p_4^2-m^2)\, \delta^{(4)}(p_1+p_2-p_3-p_4)\, |\mathcal{M}(s,t)|^2$$

$$= \frac{1}{4\,(2\pi)^2 \sqrt{s}\,\sqrt{s-4m^2}} \int \frac{d^3p_3}{2p_3^0}\; \Theta(p_1^0+p_2^0-p_3^0)$$

$$\delta\left[(p_1+p_2-p_3)^2-m^2\right] \quad |\mathcal{M}(s,t)|^2$$

$$\overset{\text{CMS}}{\downarrow}{=} \frac{1}{4\,(2\pi)^2 \sqrt{s}\,\sqrt{s-4m^2}} \int \frac{d|\vec{p}_3|\,\vec{p}_3^{\,2}\,d\Omega}{2p_3^0}\; \Theta(\sqrt{s}-p_3^0)$$

$$\underbrace{\delta(s-2\sqrt{s}\,p_3^0)}_{\frac{1}{2\sqrt{s}}\,\delta\left(p_3^0-\frac{\sqrt{s}}{2}\right)} \quad |\mathcal{M}(s,t)|^2$$

$$= \frac{1}{2}\,\frac{1}{32\pi^2 s\sqrt{s-4m^2}} \int \frac{dp_3^0\; p_3^0\; |\vec{p}_3|\, d\Omega_3}{p_3^0}\; \Theta(\sqrt{s}-p_3^0)$$

$$\times\; \delta\left(p_3^0-\frac{\sqrt{s}}{2}\right) |\mathcal{M}(s,t)|^2\; \Theta(p_3^0-m)$$

$$= \frac{1}{2}\,\frac{1}{32\pi^2 s\sqrt{s-4m^2}} \int d\Omega_3\, |\mathcal{M}(s,t)|^2 \sqrt{\frac{s}{4}-m^2}\; \Theta(\sqrt{s}-2m)$$

$$= \frac{1}{2}\,\frac{\Theta(\sqrt{s}-2m)}{64\pi^2 s} \int d\Omega_3\, |\mathcal{M}(s,t)|^2$$

<u>differential cross section</u> (identical particles in final state)



$$\sigma(p_1, p_2 \to \Delta\Omega) = \frac{1}{2} \frac{\Theta(\sqrt{s} - 2m)}{64\pi^2 s} \int_{\Delta\Omega + \Delta\bar{\Omega}} d\Omega \, |M|^2$$

$|M|^2$ symmetric under $\Theta \to \pi - \Theta$

$$\Rightarrow \sigma(p_1, p_2 \to \Delta\Omega) = \frac{\Theta(\sqrt{s} - 2m)}{64\pi^2 s} \int_{\Delta\Omega} d\Omega \, |M|^2$$

$$\frac{d\sigma}{d\Omega} = \frac{|M|^2}{64\pi^2 s} \qquad \text{(CMS)}$$

$$\sigma = \int_{\text{hemisphere}} d\Omega \, \frac{d\sigma}{d\Omega} = \frac{1}{2} \int_{\text{full sphere}} d\Omega \, \frac{d\sigma}{d\Omega}$$