

Chapter 6 : Angular Momentum

6.1. Unitary Transformations

→ We repeat general properties of the unitary operation of spatial translation

$$T_{\vec{a}} = \exp(-i\vec{a}\vec{P}/\hbar) \quad \text{which can easily cause confusion.}$$

The momentum operator \vec{P} is the generator of translations.

→ Translation acting on a wave function in \vec{x} -space :

$$\psi(\vec{x}) \rightarrow T_{\vec{a}} \psi(\vec{x}) = \exp(-i\vec{a}\vec{P}/\hbar) \psi(\vec{x}) = \exp(-i\vec{a}\vec{\nabla}) \psi(\vec{x}) = \psi(\vec{x} - \vec{a})$$

Translation acting on location eigenstates:

$$\Rightarrow \langle \vec{x} | T_{\vec{a}} \psi \rangle = \langle \vec{x} | \exp(-i\vec{a}\vec{P}/\hbar) \psi \rangle = \langle \vec{x} - \vec{a} | \psi \rangle$$

$$\Rightarrow |\vec{x}\rangle \rightarrow T_{\vec{a}} |\vec{x}\rangle = \exp(-i\vec{a}\vec{P}/\hbar) |\vec{x}\rangle = |\vec{x} + \vec{a}\rangle$$

Translation acting on momentum eigenstates:

Translation acting on momentum eigenstates:

$$|\vec{q}\rangle \rightarrow T_{\vec{a}} |\vec{q}\rangle = \exp(-i\vec{a} \vec{P}/\hbar) |\vec{q}\rangle = \exp(-i\vec{a} \vec{q}/\hbar) |\vec{q}\rangle$$

Translation acting on an operator:

Let $|q\rangle$ be a state and A a linear operator $\Rightarrow A|q\rangle$ is also a state

$$\Rightarrow T_{\vec{a}} A |q\rangle = \underbrace{T_{\vec{a}} A}_{\text{translated operator}} \underbrace{T_{-\vec{a}}}_{\text{translated state}} |\vec{q}\rangle$$

$$\hookrightarrow A \rightarrow T_{\vec{a}} A T_{-\vec{a}} = \exp(-i\vec{a} \vec{P}/\hbar) A \exp(+i\vec{a} \vec{P}/\hbar)$$

Translation acting on a function of the \vec{X} location operator: use $[X_i, P_j] = i\hbar \delta_{ij} \mathbb{1}$
 $[f(X_i), P_j] = i\hbar \delta_{ij} f'(X_i)$

$$f(\vec{X}) \rightarrow \exp(-i\vec{a} \vec{P}/\hbar) f(\vec{X}) \exp(+i\vec{a} \vec{P}/\hbar) = f(\vec{X} - \vec{a} \mathbb{1})$$

\rightarrow There is also a translation in momentum space that is generated by the $(-\vec{X})$ operator.

$$T_p^{\text{mom}} = \exp(i\vec{p} \vec{X}/\hbar)$$

This can be seen from the fact that $[f(P_i), X_j] = -i\hbar f'(P_i) \delta_{ij}$ or from the momentum space representation of the \vec{X} operator:

$$(\vec{X})_{p\text{-space}} = +i\hbar \frac{\partial}{\partial \vec{p}}$$

06/12/2017 (2)

$$\text{So we have: } |\tilde{\psi}(\vec{p})\rangle \rightarrow \exp(i\vec{q}\vec{X}/\hbar) |\tilde{\psi}(\vec{p})\rangle = |\tilde{\psi}(\vec{p}-\vec{q})\rangle$$

$$|\tilde{p}\rangle \rightarrow \exp(i\vec{q}\vec{X}/\hbar) |\tilde{p}\rangle = |\tilde{p}+\vec{q}\rangle$$

$$|\tilde{x}\rangle \rightarrow \exp(i\vec{q}\vec{X}/\hbar) |\tilde{x}\rangle = \exp(i\vec{q}\vec{x}/\hbar) |\tilde{x}\rangle$$

$$A \rightarrow \exp(i\vec{q}\vec{X}/\hbar) A \exp(-i\vec{q}\vec{X}/\hbar)$$

$$f(\vec{p}) \rightarrow \exp(i\vec{q}\vec{X}/\hbar) f(\vec{p}) \exp(-i\vec{q}\vec{X}/\hbar) = f(\vec{p}-\vec{q} \mathbb{1})$$

6.2. Orbital Angular Momentum and Spatial Rotations

→ Orbital angular momentum operator: $\vec{L} = \vec{X} \times \vec{P}$, $L_k = \epsilon_{klm} X_l P_m$ → generator for rotations

There is no issue with ordering of \vec{X} and \vec{P} operators because $\epsilon_{klm} X_l P_m = \epsilon_{klm} P_m X_l$ due to $l+m$.

→ Unitary operator for spatial rotation by angle $\tilde{\alpha}$ around axis $\frac{\vec{z}}{|\vec{z}|}$: $\exp(-i\tilde{\alpha} \vec{L}/\hbar)$

Check by an infinitesimal rotation: $\exp(-i\tilde{\alpha} \vec{L}/\hbar) = \mathbb{1} - i\tilde{\alpha} \vec{L}/\hbar \quad |\tilde{\alpha}| \ll 1$

$$\hookrightarrow (\mathbb{1} - i\tilde{\alpha} \vec{L}/\hbar) |\tilde{x}\rangle = (\mathbb{1} - i\tilde{\alpha} \epsilon_{lkm} X_l P_m / \hbar) |\tilde{x}\rangle \stackrel{m \neq l}{=} (\mathbb{1} - i\tilde{\alpha} \epsilon_{lkm} P_m X_l / \hbar) |\tilde{x}\rangle$$

Check by an infinitesimal rotation: $\exp(-i\vec{\varepsilon} \vec{L}/\hbar) = \mathbb{1} - i\vec{\varepsilon} \vec{L}/\hbar$ $|\vec{\varepsilon}| \ll 1$

$$\begin{aligned}\hookrightarrow (\mathbb{1} - i\vec{\varepsilon} \vec{L}/\hbar) |\vec{x}\rangle &= (\mathbb{1} - i\sum_{k,m} \epsilon_{k,m} X_k P_m / \hbar) |\vec{x}\rangle \stackrel{m \neq l}{=} (\mathbb{1} - i\sum_{k,m} \epsilon_{k,m} P_m X_k / \hbar) |\vec{x}\rangle \\ &= (\mathbb{1} - i\sum_{k,m} \epsilon_{k,m} P_n X_n / \hbar) |\vec{x}\rangle = (\mathbb{1} - i(\vec{\varepsilon} \times \vec{x}) \vec{P} / \hbar) |\vec{x}\rangle \\ &= \exp(-i(\vec{\varepsilon} \times \vec{x}) \vec{P} / \hbar) |\vec{x}\rangle \\ \text{Ch. 6.1.} \quad &= |\vec{x} + \vec{\varepsilon} \times \vec{x}\rangle \quad \leftarrow \text{indeed correctly rotated state (Ch. S.2)}\end{aligned}$$

$$\Rightarrow \exp(-i\vec{\varepsilon} \vec{L}/\hbar) |\vec{x}\rangle = |\mathcal{R}(\vec{\varepsilon}) \vec{x}\rangle, \mathcal{R}(\vec{\varepsilon}) : \text{spatial rotation matrix, see Ch. S.2}$$

The same should also happen to the $|\vec{p}\rangle$ state because a rotation treats all vectors equally:

$$\begin{aligned}(\mathbb{1} - i\vec{\varepsilon} \vec{L}/\hbar) |\vec{p}\rangle &= (\mathbb{1} - i\sum_{k,m} \epsilon_{k,m} X_k P_m / \hbar) |\vec{p}\rangle = (\mathbb{1} - i\sum_{k,m} \epsilon_{k,m} X_k P_m / \hbar) |\vec{p}\rangle \\ &= (\mathbb{1} + i(\vec{\varepsilon} \times \vec{p}) \vec{X} / \hbar) |\vec{p}\rangle = \exp(i(\vec{\varepsilon} \times \vec{p}) \vec{X} / \hbar) |\vec{p}\rangle \\ \text{Ch. 6.1.} \quad &= |\vec{p} + \vec{\varepsilon} \times \vec{p}\rangle \quad \leftarrow \text{yes, works!}\end{aligned}$$

$$\Rightarrow \exp(-i\vec{\varepsilon} \vec{L}/\hbar) |\vec{p}\rangle = |\mathcal{R}(\vec{\varepsilon}) \vec{p}\rangle, \mathcal{R}(\vec{\varepsilon}) : \text{spatial rotation matrix}$$

The rotation matrices $\mathcal{R}(\vec{\varepsilon})$ form the group $SO(3)$, which are the set of orthogonal 3×3 real matrices with determinant 1.

$$\hookrightarrow \mathcal{R}^T(\vec{\varepsilon}) = \mathcal{R}^{-1}(\vec{\varepsilon})$$

06/12/2017 (3)

We also check the action of the rotation operator on the \vec{X} and \vec{P} operators:

$$\begin{aligned}
 (1 - i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) \vec{X} (1 + i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) &= (1 - i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) \vec{X} (1 + i\sum_k \epsilon_{k\text{chem}} X_k P_k/\hbar) \quad \text{use } [X_i, P_m] = i\hbar \delta_{im} \mathbf{1} \\
 &\quad \downarrow \mathbf{1} + O(\epsilon^2) \qquad \qquad \downarrow O(\epsilon^2) \\
 &= (1 - i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) (1 + i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) \vec{X} + (1 - i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) (-\vec{\epsilon} \times \vec{X}) \quad \leftarrow \text{neglect } O(\epsilon^2) \text{ terms} \\
 &= \vec{X} - \vec{\epsilon} \times \vec{X} \quad \leftarrow \text{rotation in negative direction}
 \end{aligned}$$

$$\Rightarrow \exp(-i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) \vec{X} \exp(i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) = R^{-1}(\vec{\epsilon}) \vec{X}$$

$$\exp(-i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) f(\vec{X}) \exp(i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) = f(R^{-1}(\vec{\epsilon}) \vec{X})$$

$$\exp(-i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) \vec{P} \exp(i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) = R^{-1}(\vec{\epsilon}) \vec{P}$$

$$\exp(-i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) f(\vec{P}) \exp(i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) = f(R^{-1}(\vec{\epsilon}) \vec{P})$$

$$\exp(-i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) \vec{L} \exp(i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) = R^{-1}(\vec{\epsilon}) \vec{L}$$

$$\exp(-i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) f(\vec{L}) \exp(i\frac{\vec{\epsilon}}{\hbar} \vec{L}/\omega) = f(R^{-1}(\vec{\epsilon}) \vec{L})$$

} obvious to see

} same considerations

} same considerations

This makes sense since e.g. $f(R^{-1}(\vec{\epsilon}) \vec{X})$ means that we have to rotate \vec{X} with $R(\vec{\epsilon})$ to obtain the same result $f(\vec{X})$ as before the rotation. So the function f is indeed rotated by angle $|\vec{\epsilon}|$ around axis $\frac{\vec{\epsilon}}{|\vec{\epsilon}|}$.

\rightarrow Compare to $f(\vec{x}) \rightarrow f(\vec{x} - \vec{a})$ which means translation of f by $+\vec{a}$.

This makes sense since e.g. $f(R(\vec{z})\vec{x})$ means that we have to rotate \vec{x} with $R(\vec{z})$ to obtain the same result $f(\vec{x})$ as before the rotation. So the function f is indeed rotated by angle $|\vec{z}|$ around axis $\frac{\vec{z}}{|\vec{z}|}$.
 → Compare to $f(\vec{z}) \rightarrow f(\vec{z} + \vec{a})$ which means translation of f by $+\vec{a}$.

The above relations tell that \vec{L} generates the same kind of rotations on any vector operator.

This can also be expressed in the following commutation relations:

$$[L_h, X_e] = i\hbar \epsilon_{e h m} X_m$$

$$[L_h, P_e] = i\hbar \epsilon_{e h m} P_m$$

$$[L_h, L_e] = i\hbar \epsilon_{e h m} L_m$$

Finally we also check the action of the rotation operator on a wave function:

$$\underbrace{\langle \vec{x} | \exp(-i\vec{z}\vec{L}/\hbar) | \psi \rangle}_{\text{state rotated by } |\vec{z}| \text{ around } \frac{\vec{z}}{|\vec{z}|}} = \langle \vec{R}^{-1}(i\vec{z})\vec{x} | \psi \rangle = \psi(\vec{R}^{-1}\vec{x}) \quad \leftarrow \text{indeed function rotated by } R(\vec{z})$$

Infinitesimal form:

$$\langle \vec{x} | 1 - \frac{i}{\hbar} \vec{z} \cdot \vec{L} | \psi \rangle = \langle \vec{x} - \vec{z} \times \vec{x} | \psi \rangle = \psi(\vec{x} - \vec{z} \times \vec{x})$$

$$(\vec{z} \times \vec{x}) \cdot \vec{v} = \epsilon_{e h k} \epsilon_{e k l} x_m p_l \\ = \epsilon_{e h k} \epsilon_{e k l} x_m p_l = \vec{z} (\vec{x} \times \vec{v})$$

$$= \psi(\vec{x}) - (\vec{z} \times \vec{x}) \vec{v} \psi(\vec{x}) = \psi(\vec{x}) - \vec{z} (\vec{x} \times \vec{v}) \psi(\vec{x}) = \psi(\vec{x}) - \frac{i}{\hbar} (\vec{x} \times \underbrace{\frac{\vec{v}}{i} \vec{v}}_{= \vec{L}}) \psi(\vec{x})$$

$$\Rightarrow \langle \vec{x} | \vec{L} | \psi \rangle = \vec{x} \times \frac{\vec{v}}{i} \vec{v} \langle \vec{x} | \psi \rangle = \vec{x} \times \vec{p} \langle \vec{x} | \psi \rangle \quad \leftarrow \text{o.k.}$$

6.3. General Theory of the Angular Momentum

→ For any quantum mechanical system spatial rotations are described by the unitary operators

$\exp(-i\vec{J}/\hbar)$, where \vec{J} is the spatial angular momentum operator, which is Hermitian

$$\vec{J} = (J_x, J_y, J_z) = (J_1, J_2, J_3)$$

" $SU(2)$ structure constants"

These operators satisfy the angular momentum commutation relation $[J_k, J_\ell] = i\hbar \epsilon_{k\ell m} J_m$

It is convenient to consider the operators $T_h = J_h/\hbar$ to avoid factors of \hbar which satisfy

$$[T_h, T_\ell] = i\epsilon_{h\ell m} T_m \quad (\text{SU}(2) \text{ commutation relations})$$

→ Different realizations of the angular momentum operators \vec{J} (or \vec{T}) are called representations.

Examples: (1) Spin- $\frac{1}{2}$ systems : $T_h = \sigma_h/2$

(2) Spin-less particle in \mathbb{R}^3 : $T_h = -i\epsilon_{h\ell m} x_\ell \nabla_m$

(3) Spin- $\frac{1}{2}$ particle in \mathbb{R}^3 : $T_h = (\sigma_h/2) \otimes (-i\epsilon_{h\ell m} x_\ell \nabla_m)$ \otimes direct product

(1) and (2) are examples for irreducible representations, which are representations that cannot

(3) Spin- $\frac{1}{2}$ particle in \mathbb{R} : $T_k = (\sigma_k/2) \otimes (-i\epsilon_{kun} \times_e v_m)$ \otimes direct product

(1) and (2) are examples for **irreducible representations**, which are representations that cannot be written as a direct product of more elementary representations.

because $[T_u, T_v] = i\epsilon_{uvw} T_w$

It is possible to classify the irreducible representations by the eigenvalue of the operator $\vec{T}^2 = T_1^2 + T_2^2 + T_3^2$.

which commutes with each T_u , i.e. $[\vec{T}^2, T_u] = 0$, and one of the T_u

→ General Structure of the irreducible representations of the rotation group

We derive the general structure of irreducible representations where we use the eigenvalue of \vec{T}^2 to classify the representation and one of the T_u to label each state in the representation. (We take T_z .)

(e.g. Spin- $\frac{1}{2}$: $\vec{T}^2 = s(s+1)$ with $s=\frac{1}{2}$, T_z has eigenvalues $\pm \frac{1}{2}$)

We define: $T_{\pm} = T_x \pm iT_y$

$$\Rightarrow (T_{\pm})^* = T_{\mp} \quad (a)$$

$$[T_z, T_{\pm}] = iT_y \pm T_x = \pm T_{\pm} \quad (b)$$

$$[T_+, T_-] = -2i[T_x, T_y] = 2T_z \quad (c)$$

$$[\vec{T}^2, T_{\pm}] = 0 \quad (d)$$

$$\begin{aligned} T_+ T_- &= T_x^2 + T_y^2 - i[T_x, T_y] = T_x^2 + T_y^2 + T_z \\ \text{↳ } \vec{T}^2 &= T_x^2 + T_y^2 + T_z^2 = T_+ T_- - T_z + T_z^2 = T_- T_+ + T_z + T_z^2 \end{aligned} \quad (e) \quad (f)$$

06/12/2017 (5)

Step 1: The eigenvalues of \vec{T}^2 are ≥ 0 , which we therefore can write as $j(j+1)$, $j \geq 0$

On the eigenspace of \vec{T}^2 to the eigenvalue $j(j+1)$ the eigenvalues m of T_z satisfy $m^2 \leq j(j+1)$.

We can use j and m to label each angular momentum state. So we define states $|4_{jm}\rangle$ with

$$\vec{T}^2 |4_{jm}\rangle = j(j+1) |4_{jm}\rangle , \quad T_z |4_{jm}\rangle = m |4_{jm}\rangle , \quad \langle 4_{jm} | 4_{jm} \rangle = 1$$

For all states $|4\rangle$: $\underbrace{\langle 4 | \vec{T}^2 | 4 \rangle}_{\geq 0} = \underbrace{\langle 4 | T_x^2 | 4 \rangle}_{\geq 0} + \underbrace{\langle 4 | T_y^2 | 4 \rangle}_{\geq 0} + \underbrace{\langle 4 | T_z^2 | 4 \rangle}_{m^2} \geq 0$ (also true to eigenstate) ✓
 with $\langle 4 | 4 \rangle = 1$:

Step 2: The eigenspace of \vec{T}^2 to the eigenvalue $j(j+1)$ is closed w.r.t. to the operators T_x, T_y, T_z .
 So the eigenspace is also closed w.r.t. to rotations, which are functions of T_x, T_y, T_z .

Let $|4\rangle$ be an eigenstate to \vec{T}^2 with eigenvalue $j(j+1)$, then $\vec{T}^2 T_k |4\rangle = T_k \vec{T}^2 |4\rangle = j(j+1) T_k |4\rangle$ ✓

Step 3: The operators T_{\pm} raise/lower the eigenvalue of T_z by one unit

We have

It is possible to define
phase factors that way
↔

$$T_{\pm} |4_{jm}\rangle = \sqrt{j(j+1) - m(m \pm 1)} |4_{j,m \pm 1}\rangle$$

$$T_z T_{\pm} |4_{jm}\rangle \stackrel{(b)}{=} \pm T_{\pm} |4_{jm}\rangle + T_{\pm} T_z |4_{jm}\rangle = (m \pm 1) T_{\pm} |4_{jm}\rangle \quad \text{--}$$

$$\langle T_{\pm} | 4_{jm} | T_{\pm} | 4_{jm} \rangle = \langle 4_{jm} | T_{\mp} T_z | 4_{jm} \rangle$$

$$\stackrel{(c)}{=} \langle 4_{jm} | (\vec{T}^2 - T_z^2 \mp T_z) | 4_{jm} \rangle = j(j+1) - m(m \pm 1) \quad \text{--}$$

Step 4: The maximal size of eigenvalues m on a j -eigenspace is j . $|m| = j$

Step 4: The maximal size of eigenvalues m on a j -eigenspace is j : $|m| = j$

Let $m_{\max/min}$ be the largest/smallest eigenvalue of T_z on the j -eigenspace, the relation

$$T_{\pm} |4j, m_{\max/min}\rangle = \sqrt{j(j+1) - m_{\max/min}(m_{\max/min} \pm 1)} |4j, m_{\max/min} \pm 1\rangle$$

does lead to a contradiction unless $m_{\max} = j$ and $m_{\min} = -j$.

Step 5: The results from steps 1-4 are only consistent if

$$j = 0, 1, 2, \dots \quad (\text{integer spin})$$

or

$$j = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots \quad (\text{half integer spin})$$

$$\text{and } m = \underbrace{-j, -j+1, \dots, j-1, j}_{2j+1 \text{ values}}$$

For each allowed value of j the $2j+1$ orthonormal states $|4j, m\rangle$ ($m = -j, \dots, j$) form the basis of a subspace of \mathcal{H} that is invariant under rotations and represents a system with spin $-j$.

Each spin- j system forms an irreducible representation of the $SL(2)$ rotation group

For the angular momentum operators \vec{J} we have: $|4j, m\rangle = |j, m\rangle$

$$\vec{J}^2 |j, m\rangle = \hbar^2 j(j+1) |j, m\rangle, \quad J_z |j, m\rangle = \hbar m |j, m\rangle$$

$$J_{\pm} |j, m\rangle = \hbar \sqrt{j(j+1) - m(m \pm 1)} |j, m \pm 1\rangle$$

6.4. Examples for Irreducible Representations

→ Spin-0 (Trivial) Representation

$$j=0, T_h = 0$$

→ Particles without spin: e.g. Higgsboson, Mesons (Pions, kaons, ...)

→ Spin- $\frac{1}{2}$ Representation → "fundamental representation" of $Sl(2)$

$$j = \frac{1}{2}, T_h = \frac{1}{2} \tau_h, m = \frac{1}{2}, -\frac{1}{2}$$

$$\vec{T}^2 = \frac{3}{4} \mathbb{1} = \frac{1}{2} (\frac{1}{2}+1) \mathbb{1}$$

$$T_3 = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} \end{pmatrix} \quad T_+ = T_1 + i T_2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \quad T_- = T_1 - i T_2 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

$$|\frac{1}{2}, \frac{1}{2}\rangle \leftrightarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad T_3 |\frac{1}{2}, \frac{1}{2}\rangle = \frac{1}{2} |\frac{1}{2}, \frac{1}{2}\rangle \quad T_+ |\frac{1}{2}, \frac{1}{2}\rangle = 0 \quad T_- |\frac{1}{2}, \frac{1}{2}\rangle = |\frac{1}{2}, -\frac{1}{2}\rangle$$

$$|\frac{1}{2}, -\frac{1}{2}\rangle \leftrightarrow \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad T_3 |\frac{1}{2}, -\frac{1}{2}\rangle = -\frac{1}{2} |\frac{1}{2}, -\frac{1}{2}\rangle \quad T_+ |\frac{1}{2}, -\frac{1}{2}\rangle = |\frac{1}{2}, \frac{1}{2}\rangle \quad T_- |\frac{1}{2}, -\frac{1}{2}\rangle = 0$$

→ Spin-1 Representation → 3-dimensional representation, "fundamental representation" of $SO(3)$

$$j = 1, \vec{T}^2 = 2 \mathbb{1} = 1 (1+1) \mathbb{1}, m = +1, 0, -1$$

Spin basis: $\{|1, m\rangle\} = \{|1, +1\rangle, |1, 0\rangle, |1, -1\rangle\}$ with $T_3 |1, m\rangle = m |1, m\rangle$

$$j=1, \quad l=\langle 1 \rangle = 1(1+1)\frac{1}{2}, \quad m=+1, 0, -1$$

Spin basis: $\{|1,m\rangle\} = \{|1,+1\rangle, |1,0\rangle, |1,-1\rangle\}$ with $T_3|1,m\rangle = m|1,m\rangle$

$$T_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad T_+ = \begin{pmatrix} 0 & \sqrt{2} & 0 \\ 0 & 0 & \sqrt{2} \\ 0 & 0 & 0 \end{pmatrix} \quad T_- = \begin{pmatrix} 0 & 0 & 0 \\ -\sqrt{2} & 0 & 0 \\ 0 & -\sqrt{2} & 0 \end{pmatrix} \quad |1,1\rangle \leftrightarrow \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, |1,0\rangle \leftrightarrow \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, |1,-1\rangle \leftrightarrow \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

$$T_1 = \frac{1}{2}(T_+ + T_-) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \quad T_2 = \frac{1}{2i}(T_+ - T_-) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -i & 0 \\ i & 0 & -i \\ 0 & i & 0 \end{pmatrix}$$

The spin basis is useful for the description of spin-1 particles in analogy to the description of spin- $\frac{1}{2}$.

In the spin basis the 3 basis states $|1,m\rangle$ ($m=1, 0, -1$) do not describe the 3 spatial directions (x, y, z) . However, there is an alternative basis where the 3 basis states indeed correspond to the spatial directions. This basis is called the "adjoint representation of $SU(2)$ ".

Adjoint Representation:

The adjoint representation is obtained from the $SU(2)$ structure constants and can be derived from the infinitesimal rotations of spatial 3-vectors.

generators of the adjoint rep

↪ Rotation by $|\vec{\varepsilon}| \ll 1$ around axis $\frac{\vec{\varepsilon}}{|\vec{\varepsilon}|}$: $\vec{x} \rightarrow \vec{x} + \vec{\varepsilon} \times \vec{x} = (\mathbb{1} - i \frac{\vec{\varepsilon}}{|\vec{\varepsilon}|} \vec{t}) \vec{x}$

$$= \underbrace{(t_{nk})}_e$$

$$x_n \rightarrow x_n + \epsilon_{nkl} \varepsilon_l x_k = (\delta_{nk} - i \varepsilon_k [-i \epsilon_{nkl}]) x_k$$

$$\Rightarrow t_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix} \quad t_2 = \begin{pmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ -i & 0 & 0 \end{pmatrix} \quad t_3 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

06/12/2017 (7)

One can show that: $T_k = U t_k U^+$ with $U = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 & i & 0 \\ 0 & 0 & 1 \\ 1 & i & 0 \end{pmatrix}$

This means that a spatial 3-vector $\vec{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$ has the form $U \vec{x} = \frac{1}{\sqrt{2}} \begin{pmatrix} -x+iy \\ z \\ x+iy \end{pmatrix}$ in the spin-1 basis.

→ Comment:

All representations of angular momentum with half-integer j (i.e. $j = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots$) are not representations of the spatial rotation group $SO(3)$. This is only possible for integer j (i.e. $j = 0, 1, 2, \dots$).


6.5. Spherical harmonic functions

→ The spherical harmonic functions are the angular momentum analogue to the eigenfunctions $\langle \vec{x} | \vec{p} \rangle = \frac{e^{i\vec{p}\cdot\vec{x}}}{(2\pi)^{3/2}}$ of the momentum operator \vec{P} .

The spherical harmonic functions are an explicit realization of the irreducible representations of angular momentum and the rotation group for all integer values of j , i.e. for $j=0, 1, 2, \dots$

→ We have to switch from cartesian to spherical (polar) coordinates:

$$x = x_r = r \sin\theta \cos\phi, \quad y = x_\theta = r \sin\theta \sin\phi, \quad z = x_\phi = r \cos\theta$$



→ We have to switch from Cartesian to Spherical (polar) coordinates:

$$x = x_1 = r \sin\theta \cos\phi, \quad y = x_2 = r \sin\theta \sin\phi, \quad z = x_3 = r \cos\theta$$

In spherical coordinates the basis vectors at each point $\vec{x} = r \vec{e}_r$ in space depend on the $(r, \theta, \phi) = (\text{radius}, \text{polar angle}, \text{azimuthal angle})$ values of that point \vec{x} .

$$\vec{e}_r = \vec{e}_x \sin\theta \cos\phi + \vec{e}_y \sin\theta \sin\phi + \vec{e}_z \cos\theta$$

$$\theta \rightarrow \theta + \frac{\pi}{2} \quad \theta = \frac{\pi}{2}, \quad \phi \rightarrow \phi + \frac{\pi}{2}$$

$$\vec{e}_\theta = \vec{e}_x \cos\theta \cos\phi + \vec{e}_y \cos\theta \sin\phi - \vec{e}_z \sin\theta$$

$$\vec{e}_\phi = -\vec{e}_x \sin\phi + \vec{e}_y \cos\phi$$

Form of the nabla: $\vec{\nabla} = \vec{e}_r \frac{\partial}{\partial r} + \vec{e}_\theta \frac{1}{r} \frac{\partial}{\partial \theta} + \vec{e}_\phi \frac{1}{r \sin\theta} \frac{\partial}{\partial \phi}$

$$\vec{e}_r \times \vec{e}_\theta = \vec{e}_\phi$$

$$\vec{e}_r \times \vec{e}_\phi = -\vec{e}_\theta$$

Angular momentum operator:

$$\vec{L} = \vec{x} \times \frac{i}{i} \vec{\nabla} = \frac{i}{i} r \vec{e}_r \times \vec{\nabla} = \frac{i}{i} \left(\vec{e}_\phi \frac{\partial}{\partial \theta} - \vec{e}_\theta \frac{1}{\sin\theta} \frac{\partial}{\partial \phi} \right)$$


$$L_x = L_1 = \frac{i}{i} \left(-\sin\phi \frac{\partial}{\partial \theta} - \cos\phi \cot\theta \frac{\partial}{\partial \phi} \right)$$

$$L_y = L_2 = \frac{i}{i} \left(\cos\phi \frac{\partial}{\partial \theta} - \sin\phi \cot\theta \frac{\partial}{\partial \phi} \right)$$

$$L_z = L_3 = \frac{i}{i} \frac{\partial}{\partial \phi}$$

$$\left. \begin{aligned} L_\pm &= i e^{\pm i \phi} \left(\pm \frac{\partial}{\partial \theta} + i \cot\theta \frac{\partial}{\partial \phi} \right) \\ \vec{L}^2 &= -i^2 \left[\frac{1}{\sin\theta} \frac{\partial}{\partial \theta} \left(\sin\theta \frac{\partial}{\partial \theta} \right) + \frac{1}{\sin^2\theta} \frac{\partial^2}{\partial \phi^2} \right] \end{aligned} \right\}$$

$$d\Omega = d\cos\theta d\phi = \sin\theta d\theta d\phi$$



06/12/2017 (8)

→ The angular momentum operators L_L only act on the polar and azimuthal angles.
So they act on functions $Y(\theta, \phi)$ defined on the surface of the unit sphere (called S^2)

The set of complex-valued functions defined on the surface of the unit sphere

$$L^2(S^2) = \left\{ Y(\theta, \phi) : S^2 \rightarrow \mathbb{C} \mid \int d\Omega |Y(\theta, \phi)|^2 = \int_0^\pi d\theta \sin\theta \int_0^{2\pi} d\phi |Y(\theta, \phi)|^2 < \infty \right\}$$

with the scalar product $\langle Y_1, Y_2 \rangle = \int_{S^2} d\Omega Y_1^*(\Omega) Y_2(\Omega)$, $\Omega = (\theta, \phi)$ is a Hilbert space.

→ The spherical harmonic functions $Y_{lm}(\theta, \phi) \in L^2(S^2)$ are the simultaneous eigenfunctions of the operators \vec{L}^2 and L_z with

$$\vec{L}^2 Y_{lm}(\Omega) = l(l+1) Y_{lm}(\Omega), \quad l = 0, 1, 2, \dots$$

$$L_z Y_{lm}(\theta, \phi) = m Y_{lm}(\theta, \phi), \quad m = -l, -l+1, \dots, l-1, l$$

and they form a CONS of $L^2(S^2)$.

We determine the $Y_{lm}(\theta, \phi)$ by using the separation ansatz $Y_{lm}(\theta, \phi) = P_{lm}(\cos \theta) b_m(\phi)$ and starting with the eigenvalue equation

$$L_z Y_{lm} = m Y_{lm} \iff \frac{\partial}{\partial \phi} b_m(\phi) = i m b_m(\phi) \implies b_m(\phi) = e^{im\phi}$$

We determine the $Y_{lm}(\theta, \phi)$ by using the separation ansatz $Y_{lm}(\theta, \phi) = P_{lm}(\cos \theta) b_m(\phi)$ and starting with the eigenvalue equation

$$L_2 Y_{lm} = \text{tr}_m Y_{lm} \iff \frac{\partial^2}{\partial \phi^2} b_m(\phi) = i m b_m(\phi) \Rightarrow b_m(\phi) = e^{im\phi}$$

Thus the second eigenvalue equation $L^2 Y_{lm} = h^2 l(l+1) Y_{lm}$ can be rewritten as

$$\left[\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) - \frac{m^2}{\sin^2 \theta} + l(l+1) \right] P_{lm}(\cos \theta) = 0$$

We change variables : $\xi = \cos \theta \rightarrow \frac{\partial}{\partial \theta} = \frac{d \cos \theta}{d \theta} \frac{\partial}{\partial \xi} = - \sin \theta \frac{\partial}{\partial \xi} \quad (-1 \leq \xi \leq 1)$

$$\Rightarrow \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) = \frac{\partial}{\partial \xi} \left(\sin^2 \theta \frac{\partial}{\partial \xi} \right) = \frac{\partial}{\partial \xi} \left((1-\xi^2) \frac{\partial}{\partial \xi} \right) = (1-\xi^2) \frac{\partial^2}{\partial \xi^2} - 2\xi \frac{\partial}{\partial \xi}$$

$$\Rightarrow \left[(1-\xi^2) \frac{\partial^2}{\partial \xi^2} - 2\xi \frac{\partial}{\partial \xi} + l(l+1) - \frac{m^2}{1-\xi^2} \right] P_{lm}(\xi) = 0$$

This is the defining equation of the associated Legendre polynomials.

The P_{lm} can be obtained from the (regular) Legendre polynomials $P_l(\xi) = \frac{1}{2^l l!} \frac{d^l}{d\xi^l} (\xi^2 - 1)^l$, $l=0,1,\dots$ by the relation

$$P_{lm}(\xi) = (1-\xi^2)^{\frac{m}{2}} \frac{d^m}{d\xi^m} P_l(\xi) \quad \text{with } m \geq 0.$$

The regular Legendre polynomials form a complete set of orthogonal functions on the interval $[-1, 1]$ with

$$\int_{-1}^1 d\xi P_n(\xi) P_m(\xi) = \frac{2}{2m+1} \delta_{nm}$$

They satisfy the recursion relations

$$(l+1) P_{l+1}(\xi) = (2l+1)\xi P_l(\xi) - l P_{l-1}(\xi), \quad (1-\xi^2) \frac{d}{d\xi} P_l(\xi) = l (P_{l-1}(\xi) - \xi P_l(\xi))$$

Lowest Legendre polynomials: $P_0(\xi) = 1, P_1(\xi) = \xi, P_2(\xi) = \frac{1}{2}(3\xi^2 - 1), P_3(\xi) = \frac{1}{2}(5\xi^3 - 3\xi), \dots$

06/12/2017 (9)

The associated Legendre polynomials have the properties

$$\int_{-1}^1 d\xi P_{\ell m}(\xi) P_{\ell' m'}(\xi) = \frac{2}{2\ell+1} \frac{(\ell+m)!}{(\ell-m)!} \delta_{\ell\ell'} \quad (m \geq 0)$$

$$P_{\ell m}(-\xi) = (-1)^{\ell+m} P_{\ell m}(\xi)$$

$$P_{\ell 0}(\xi) = P_\ell(\xi) \quad , \quad P_{\ell 0}(\xi) = (2\ell-1)!! \quad (1-\xi^2)^{\frac{\ell}{2}}$$

$$(2\ell-1)!! := (2\ell-1)(2\ell-3)\dots 3 \cdot 1$$

→ Explicit form of the spherical harmonic functions:

$$Y_{\ell m}(\theta, \phi) = (-1)^{\frac{(m+l+m)}{2}} \left[\frac{2\ell+1}{4\pi} \frac{(\ell-l+m)!}{(\ell+l+m)!} \right]^{\frac{1}{2}} P_{\ell m}(\cos\theta) e^{im\phi}$$

$$\text{Orthonormality: } \int d\Omega Y_{\ell m}^*(\Omega) Y_{\ell' m'}(\Omega) = \delta_{\ell\ell'} \delta_{mm'}$$

$$\text{Completeness: } \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} Y_{\ell m}(\theta, \phi) Y_{\ell m}^*(\theta', \phi') = \delta(\cos\theta - \cos\theta') \delta(\phi - \phi') = \frac{1}{\sin\theta} \delta(\theta - \theta') \delta(\phi - \phi')$$

$$\text{Addition theorem: } \sum_{m=-\ell}^{\ell} Y_{\ell m}(\theta, \phi) Y_{\ell m}(\theta', \phi') = \frac{2\ell+1}{4\pi} P_\ell(\cos\alpha), \quad \alpha : \text{angle between directions } (\theta, \phi) \text{ and } (\theta', \phi')$$

$$\cos\alpha = \cos\theta \cos\theta' + \sin\theta \sin\theta' \cos(\phi - \phi')$$

$$\text{Parity transformation: } \vec{x} \rightarrow P\vec{x} = -\vec{x} \Rightarrow (\theta, \phi) \rightarrow (\pi - \theta, \phi + \pi)$$

$$P Y_{\ell m}(\theta, \phi) = Y_{\ell m}(\pi - \theta, \phi + \pi) = (-1)^\ell Y_{\ell m}(\theta, \phi)$$

$$\cos(\pi - \theta) = -\cos\theta$$

Parity transformation: $\vec{r} \rightarrow -\vec{r}$ $\Rightarrow (\theta, \phi) \rightarrow (\pi - \theta, \phi + \pi)$

$$P Y_{lm}(\theta, \phi) = Y_{lm}(\pi - \theta, \phi + \pi) = (-1)^l Y_{lm}(\theta, \phi)$$

$$\cos(\pi - \theta) = -\cos \theta$$

$$m\text{-Symmetry: } Y_{-lm}(\theta, \phi) = (-1)^m Y_{lm}^*(\theta, \phi)$$

Some explicit expressions: (use m -symmetry to obtain expressions for $m < 0$)

$$Y_{00} = \frac{1}{\sqrt{4\pi}}$$

$$Y_{11} = -\sqrt{\frac{3}{8\pi}} \sin \theta e^{i\phi}, \quad Y_{10} = \sqrt{\frac{3}{4\pi}} \cos \theta$$

$$Y_{22} = \sqrt{\frac{15}{32\pi}} \sin^2 \theta e^{2i\phi}, \quad Y_{21} = -\sqrt{\frac{15}{8\pi}} \sin \theta \cos \theta e^{i\phi}, \quad Y_{20} = \sqrt{\frac{5}{16\pi}} (3 \cos^2 \theta - 1)$$


→ The completeness property of the spherical harmonic functions Y_{lm} allows to write every wave function $\psi(\vec{r}) \in L^2(\mathbb{R}^3)$ as

$$\psi(\vec{r}) = \psi(r, \theta, \phi) = \sum_{l=0}^{\infty} \sum_{m=-l}^{+l} u_{lm}(r) Y_{lm}(\theta, \phi) \quad \text{where}$$

$$u_{lm}(r) = \int_0^\pi d\theta \sin \theta \int_0^{2\pi} d\phi Y_{lm}^*(\theta, \phi) \psi(\vec{r}) = \int_0^\pi d\cos \theta \int_0^{2\pi} d\phi Y_{lm}^*(\theta, \phi) \psi(\vec{r})$$

angular / multipole spectrum

06/12/2017 (10)




Schuskel p. 119

Abb. 5.6. Polardiagramme der Bahndrehimpulseigenfunktionen Y_{lm} mit $l = 0, 1, 2, 3$


CMB = Cosmic Microwave
Background

What do we learn from CMB observations - Rubakov, V.A. et al. Phys.Atom.Nucl. 75 (2012) 1123-1141 arXiv:1008.1704 [astro-ph.CO] ITEP-TH-30-10

$$T = 2.725^\circ K, \frac{\delta T}{T} \sim 10^{-5}$$



~The CMB temperature map, obtained by WMAP experiment [\cite{WMAP}](#). The brighter a region is, the hotter radiation comes from it.



~The angular spectrum of the CMB temperature anisotropy [\cite{ACBAR}](#). The line is a prediction of the standard Λ CDM model. The quantity in vertical axis is D_l defined by [\(???\)](#).