

Exercises for T2, Summer term 2017, Sheet 7

1) Ladder operators

The ladder operators of a one-dimensional harmonic oscillator a, a^\dagger fulfill the commutation relation $[a, a^\dagger] = \mathbb{1}$. Show that:

1. $[a, (a^\dagger)^n] = n(a^\dagger)^{n-1}$
2. $[a, f(a^\dagger)] = f'(a^\dagger)$

Assume that the function f is defined as a power series.

2) Commutators

Determine the following commutators:

In one dimension: $[X^2, P^2]$

In three dimensions: The operator A is defined as $A = X_1P_2 - X_2P_1$. Determine $[A, X_1^2]$.

3) Expectation values for the harmonic oscillator

Calculate the expectation values $\langle X \rangle_n, \langle P \rangle_n, \langle X^2 \rangle_n, \langle P^2 \rangle_n$ for the energy eigenstates $|n\rangle$ of the harmonic oscillator ($\langle O \rangle_n \equiv \langle n|O|n\rangle$). Use algebraic methods with the ladder operators.

4) Coherent state I

The state $|z\rangle \equiv |\psi_z\rangle$ (with $z \in \mathbb{C}$) of a harmonic oscillator is defined via the eigenvalue equation $a|z\rangle = z|z\rangle$. Show that the solution of this equation is

$$|z\rangle = C e^{za^\dagger} |0\rangle.$$

Use the equations derived in exercise (1). The state $|z\rangle$ is an example of a coherent state.

5) Coherent state II

Write the coherent state $|z\rangle$ as a linear combination of the normalized energy eigenstates $|n\rangle \equiv |\phi_n\rangle$ of the harmonic oscillator. Calculate $\langle n|z\rangle$. Determine the normalization constant C (up to an arbitrary phase $e^{i\alpha}$) by imposing the normalization condition $\langle z|z\rangle = 1$.